

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 November 2001 (22.11.2001)

PCT

(10) International Publication Number
WO 01/87849 A2

(51) International Patent Classification⁷: **C07D 233/54**,
A61K 31/395, A61P 43/00, C07D 401/04, 401/14, 405/12,
409/12, 403/12

(21) International Application Number: **PCT/US01/15027**

(22) International Filing Date: **10 May 2001 (10.05.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/203,784 12 May 2000 (12.05.2000) **US**
60/205,213 18 May 2000 (18.05.2000) **US**

(71) Applicant (for all designated States except US): **GEN-
ZYME CORPORATION** [US/US]; One Kendall Square,
Cambridge, MA 02139 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SNEDDON, Scott,**
F. [US/US]; 38 Washington Square South, Salem, MA
01970 (US). **KANE, John, L.** [US/US]; 10 O'Moore
Avenue, Maynard, MA 01754 (US). **HIRTH, Bradford,**

H. [US/US]; 4 Dogwood Road, Littleton, MA 01460 (US).
VINICK, Fred [US/US]; 5 Litchfield Road, Lexington,
MA 02420 (US). **QIAO, Shuang** [CN/US]; 334 Harvard
Street, Apt. J-3, Cambridge, MA 02139 (US). **NAHILL,**
Sharon, R. [US/US]; 21 Vincent Avenue, Belmont, MA
02478 (US).




(74) Agents: **BROOK, David, E.** et al.; Hamilton, Brook,
Smith & Reynolds, P.C., 530 Virginia Road, P.O. Box
9133, Concord, MA 01742-9133 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: **MODULATORS OF TNF- α SIGNALING**



(57) Abstract: The present invention provides compounds which are modulators of TNF- α signaling and methods of use thereof for treating a patient having a TNF- α mediated condition. The compounds can be represented by structural formulas (I), (II) and (III).



Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

MODULATORS OF TNF- α SIGNALING

RELATED APPLICATIONS

This application claims the benefit of US Provisional Application No. 60/203,784, filed May 12, 2000, and US Provisional Application No. 60/205,213, filed May 18, 2000. The entire teachings of these applications are incorporated herein by reference.

BACKGROUND OF THE INVENTION

The cytokine tumor necrosis factor α ("TNF- α ") has a broad spectrum of biological activities. TNF- α is produced by activated macrophages and a variety of other cells, including antigen-stimulated T cells, activated mast cells and activated natural killer cells. TNF- α is initially produced as a transmembrane protein of about 25 kD. A 17 kD fragment of this membrane protein is proteolytically cleaved from the cell membrane and circulates as a 51 kD homotrimer. TNF- α mediated processes proceed via the interaction of this trimeric protein with a receptor protein at the surface of a target cell.

TNF- α plays an important role in coordinating the body's response to infection, and serves as an important mediator of inflammation. For example, TNF- α signaling has been implicated in the induction of fever and the production of interferon- γ by T cells. TNF- α induces increased binding of leukocytes to endothelial cells, resulting in accumulation of leukocytes at sites of infection. TNF- α signaling has also been implicated in inducing the production of interleukin-1 and prostaglandins by macrophages, and is involved in the breakdown of the extracellular matrix, inducing collagenase in synoviocytes, and in bone resorption via osteoclast activation.

TNF- α has certain effects on the growth and metastatic potential of tumors. For example, certain human tumor cell lines are sensitive to TNF- α *in vitro* and TNF- α activation may precede killing of tumor cells by macrophages.

High levels of TNF- α are generally associated with chronic immune or inflammatory diseases, and are considered a cause of neural and cellular degeneration. At lower levels, however, TNF- α plays an important role in the cell

-2-

life cycle, cellular response to foreign attack, and maintenance of homeostasis. For this reason, it will be appreciated that the purpose of this invention is not the complete and absolute inhibition of TNF- α , but rather the modulation of the cellular response to TNF- α levels and the treatment of TNF- α mediated conditions, thereby
5 permitting an effective treatment for the chronic immune and inflammatory responses that occur when excess TNF- α is produced.


The production of TNF- α has been implicated in a variety of disease states including but not limited to the following: septic shock; endotoxic shock; cachexia syndromes associated with bacterial infections, such as tuberculosis and meningitis;
10 viral infections, such as AIDS; parasitic infections, such as malaria; neoplastic disease; autoimmune disease, including some forms of arthritis, especially rheumatoid and degenerative forms; and adverse effects associated with treatment for the prevention of graft rejection. Thus, there is a need for agents which can interrupt or modulate the TNF- α signaling process.

15 SUMMARY OF THE INVENTION

The present invention provides compounds which are modulators of TNF- α signaling and methods of use thereof for treating a patient having a TNF- α mediated condition.

In one embodiment, the invention provides compounds of Formula (I),


20



-3-

In Formula (I), R_1 is H or NH_2 ; R_2 and R_3 are each, independently, -H, -OH, a substituted or unsubstituted alkyl, or a substituted or unsubstituted alkoxy; R_4 is, -H or a substituted or unsubstituted alkyl; X is O, S, CH_2 or SO_2 ; V, W and Z are each, independently, N or CH; Y is substituted and unsubstituted phenyl or a substituted and unsubstituted heterocyclyl; and n is 0, 1 or 2.


In another embodiment, the invention provides compounds of Formula II,



where R_5 is substituted or unsubstituted aralkyl, a substituted or unsubstituted cycloalkyl or a substituted or unsubstituted cycloalkylalkyl; R_6 is -H or $-NR_{13}R_{14}$; R_7 is substituted or unsubstituted phenyl; and R_{13} and R_{14} are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl or R_{13} and R_{14} together with the nitrogen to which they are attached is a heterocycloalkyl.

The present invention further relates to compounds of Formula III,

-4-



where R_8 and R_{12} are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl or a substituted or unsubstituted heteroaralkyl; R_9 is -H, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted heteroaralkyl; R_{10} is substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaralkyl or a substituted or unsubstituted heterocycloalkylalkyl; and R_{11} is substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl, a substituted or unsubstituted cycloalkylalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted heteroaralkyl, a substituted or unsubstituted benzophenonyl or a substituted or unsubstituted cycloalkylalkyl.

In yet another embodiment, the present invention relates to a method of treating a TNF- α mediated condition in a patient. The method comprises the step of administering to the patient a therapeutically effective amount of at least one compound of Formula I, Formula II or Formula III, as defined above.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to compounds which are antagonists of TNF- α signaling and, therefore, are effective agents for the treatment of TNF- α mediated medical conditions, disorders and diseases, such as chronic inflammation, tissue breakdown and cancer.

For the purposes of the present invention, the language "alkyl" is intended to include a straight chain or branched saturated hydrocarbyl group. Preferred alkyl groups include C_1 - C_{12} -alkyl groups, while more preferred alkyl groups include C_1 - C_6 -alkyl groups. The language "cycloalkyl" is intended to include a mono-, bi- or

polycyclic alkyl group. Preferred cycloalkyl groups include monocyclic C₃-C₈-cycloalkyl groups. The language "alkoxy" is intended to include an alkyl-O- group or a cycloalkyl-O- group, where the preferred alkyl and cycloalkyl groups are those given above. The language "aromatic ether" is intended to include an -O-aryl or -O-heteroaryl. The language "alkenyl" is intended to include a straight chain or branched hydrocarbyl group which includes one or more double bonds. Preferred alkenyl groups include C₂-C₁₂-alkenyl groups. The language "cycloalkenyl" is intended to include a cyclic hydrocarbyl group which includes one or more double bonds but is not aromatic. Preferred cycloalkenyl groups include C₅-C₈-cycloalkenyl groups.

The language "aryl" is intended to include an aromatic carbocyclic group, such as a phenyl group, a naphthyl group or a phenyl or naphthyl group which is fused with a five or six-membered saturated, partially unsaturated or aromatic carbocyclic ring.

The language "heterocycle" and "heterocyclic group" is intended to include a saturated, aromatic or partially unsaturated ring system which includes at least one heteroatom, such as one or more oxygen, nitrogen or sulfur atoms or a combination thereof.

The language "heterocycloalkyl" is intended to include saturated heterocyclic groups, such as piperidyl, pyrrolidyl, piperazyl, tetrahydrofuranyl and morpholyl.

The language "heteroaryl" is intended to include an aromatic heterocyclic group. Suitable heteroaryl groups include, but are not limited to, pyridyl, pyrimidyl, quinolyl, isoquinolyl, pyrrolyl, quinoxalyl, imidazolyl, oxazolyl, isoxazolyl, pyrazolyl, thienyl, furanyl, pyrazolyl, thiadiazolyl, oxadiazolyl, indazolyl, thiazolyl, isothiazolyl, and tetrazolyl. Heteroaryl groups also include ring systems in which a carbocyclic aromatic ring, carbocyclic non-aromatic ring or heteroaryl ring is fused to one or more other heteroaryl rings, *e.g.*, benzo(b)thienyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, benzoxadiazolyl, indolyl, tetrahydroindolyl, azaindolyl, indazolyl, quinoliny, imidazopyridyl, puryl, pyrrolo[2,3-d]pyrimidyl, pyrazolo[3,4-d]pyrimidyl.

The language "aralkyl" is intended to include an alkyl group which is

substituted by one or more aryl groups. A substituted aralkyl can have a substituent on the aryl or on the alkyl portion of the aralkyl. Preferred aralkyl groups include benzyl, diphenylmethyl and 2-phenethyl groups. The language "heteroaralkyl" is intended to include an alkyl group which is substituted by a heteroaryl group or by a heteroaryl group and one or more aryl groups. A substituted heteroaralkyl can have a substituent on the a heteroaryl or on the alkyl portion of the heteroaralkyl. Preferably, a heteroaryl group is an alkyl group substituted by a heteroaryl group.

The language "cycloalkylalkyl" is intended to include an alkyl group substituted with a cycloalkyl group.

10 The language "heterocycloalkylalkyl" is intended to include an alkyl group substituted with a heterocycloalkyl group.


Alkyl, cycloalkyl, alkenyl, cycloalkenyl, the alkyl portion of an aralkyl, the alkyl portion of a heteroaralkyl, cycloalkylalkyl, and alkoxy groups can be substituted or unsubstituted. Substituted groups of this type can include one or more substituents independently selected from halo, such as fluoro, chloro, bromo and iodo; alkyl, such as C₁-C₆-alkyl; nitro; hydroxyl; -NR₁₃R₁₄, wherein R₁₃ and R₁₄ are defined as above; -C(O)R₁₅, wherein R₁₅ is -H, an alkyl, an aryl or an aralkyl; cyano; aryl groups; cycloalkyl groups and heterocyclic groups, such as heteroaryl groups.

20 Aryl, heterocyclic, such as heteroaryl, aromatic ethers, the aromatic portion of an aralkyl, the aromatic portion of heteroaralkyl, heterocycloalkylalkyl and benzophenone groups can be substituted or unsubstituted. Suitable substituents include one or more substituents independently selected from halo, such as fluoro, chloro, bromo or iodo; alkyl, preferably C₁-C₃-alkyl; a halogenated alkyl, preferably a halogenated C₁-C₃-alkyl; alkoxy, preferably C₁-C₃-alkoxy; aromatic ether; alkyl substituted with an aromatic ether; a hydroxy substituted alkyl; alkoxy substituted aromatic ether; -S-(alkyl); cyano; azide; nitro; -C(O)R₁₅; -NR₁₃R₁₄; -C(O)NR₁₃R₁₄; -C(O)OR₁₆, wherein R₁₆ is -H, an alkyl, an aryl or an aralkyl; benzyl; 4-(4-benzylpiperazin-1-yl)methyl; 4-4-(2-fluorophenyl)piperazin-1-yl)methyl; a halogenated aryl; methylenedioxy; an aryl; a heteroaralkyl; a heterocycloalkylalkyl; and a heterocyclic, such as a heteroaryl group.

30

-7-

In one embodiment, the present invention provides compounds of Formula I,




In Formula (I), R_1 is H or NH_2 ; R_2 and R_3 are each, independently, -H, -OH, a substituted or unsubstituted alkyl or a substituted or unsubstituted alkoxy; R_4 is, -H or a substituted or unsubstituted alkyl; X is O, S, CH_2 or SO_2 ; V, W and Z are each, independently, N or CH; Y is substituted and unsubstituted phenyl or a substituted and unsubstituted heterocyclyl; and n is 0, 1 or 2.

In one embodiment, Y is a phenyl group which has one or more substituents independently selected from the group consisting of halogen, linear or branched C_1 - C_4 -alkoxy, trifluoromethoxy, dioxymethylene, hydroxyalkyl, trifluoromethyl, $HC(O)-$, linear or branched C_1 - C_4 -alkyl, heterocyclyl and substituted or unsubstituted heterocycloalkylalkyl. Preferred substituents for Y include fluoro, chloro, methoxy, morpholyl, N-morpholinomethyl, tetrahydroisoquinolyl, tetrahydroisoquinolinomethyl, 4-(4-benzyl-piperazin-1-yl)methyl, 4-(4-(2-fluoro-phenyl)piperazin-1-yl)methyl and isopropyl. Y can also be a heterocyclyl group, *e.g.*, pyridyl, furyl or pyrrolidyl.

-8-


Alternatively in Formula (I), R_1 - R_4 , V, W, X, Z and n are as described above, and Y is represented by the following structural formula:



wherein R_{50} and R_{51} are independently an alkyl group, a substituted alkyl group, an
 5 aryl group a substituted aryl group, or, taken together with the nitrogen atom to
 which they are bonded, are a substituted heterocycloalkyl, an unsubstituted
 heterocycloalkyl, a substituted heteroaryl group or an unsubstituted heteroaryl group.
 Preferably, R_{50} and R_{51} are, taken together with the nitrogen atom to which they are
 bonded, an *N*-substituted piperazyl group, wherein the *N*-substituent is an aryl group,
 10 a substituted aryl group, a $-\text{CH}_2$ -aryl group or a $-\text{CH}_2$ -(substituted aryl group), and
 preferably phenyl, substituted phenyl, benzyl or substituted phenyl. In a preferred
 embodiment, R_1 and R_4 are -H, R_2 - R_3 are methyl, V and Z are each -CH-, W is -N-,
 X is -O- and n is 0. Suitable substituents for a heterocycloalkyl or heteroaryl group
 formed by R_{50} and R_{51} taken together with the nitrogen atom to which they are
 15 bonded are as described below for substituted heterocycles.

In another embodiment, the invention provides compounds of Formula II,

-9-




where R_5 is substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted cycloalkylalkyl; R_6 is -H or $-NR_{13}R_{14}$; R_7 is substituted or unsubstituted phenyl; and R_{13} and R_{14} are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl or R_{13} and R_{14} together with the nitrogen to which they are attached is a heterocycloalkyl.

In one embodiment, R_5 is substituted or unsubstituted benzyl. Suitable substituents on the benzyl group include halogen atoms and linear and branched C_1 - C_4 -alkoxy. Preferably, R_5 is unsubstituted benzyl or benzyl having one or more substituents independently selected from chloro and methoxy. In another embodiment, R_5 is C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkyl- C_1 - C_4 -alkyl or substituted or unsubstituted phenyl- C_2 - C_4 -alkyl. For example, R_5 can be 2-phenethyl, cyclohexyl or cyclopentylethyl.

R_7 is, preferably, phenyl having one or more of the following independently selected substituents: halogen, linear C_1 - C_6 -alkyl, branched C_1 - C_6 -alkyl, cyclic C_3 - C_6 -alkyl or trifluoromethyl. More preferably, R_7 is phenyl having one or more of the following independently selected substituents: fluoro, chloro, and linear C_1 - C_4 -alkyl or branched C_1 - C_4 -alkyl.

The present invention further relates to compounds of Formula III



where R_8 and R_{12} are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, substituted or unsubstituted aralkyl or a substituted
 5 or unsubstituted heteroaralkyl; R_9 is -H, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl, a substituted or unsubstituted heteroaryl or a substituted or unsubstituted heteroaralkyl; R_{10} is substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaralkyl or a substituted or unsubstituted heterocycloalkylalkyl; and R_{11} is substituted or
 10 unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl, a substituted or unsubstituted cycloalkylalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted heteroaralkyl, a substituted or unsubstituted benzophenone or a substituted or unsubstituted cycloalkylalkyl.

In a preferred embodiment, one of R_8 and R_{12} is -H and the other is
 15 substituted or unsubstituted phenyl, phenyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -alkyl, linear C_1 - C_{12} -alkyl, branched C_1 - C_{12} -alkyl, cyclic C_3 - C_{12} -alkyl or dicycloalkyl- C_1 - C_4 -alkyl. Examples of suitable substituents for the phenyl group(s) of R_8 or R_{12} include one or more of the following independently selected groups: alkoxy, such as C_1 - C_4 -alkoxy, preferably methoxy; alkyl, such as C_1 - C_4 -alkyl, preferably methyl and ethyl; and
 20 cyano. Suitable identities for R_8 or R_{12} include, but are not limited to, 2,2-diphenylethyl, 2-(4-ethylphenyl)ethyl, benzyl, diphenylmethyl, 1,2-diphenylethyl, 3,3-diphenylpropyl, 3,4,5-trimethoxybenzyl, 2,4,4-trimethylisopentyl, 2-(4-methoxyphenyl)ethyl, 2-cyclopentyl-2-phenylethyl, or 2-phenyl-2-pyridylethyl.

R_9 is, preferably, substituted or unsubstituted phenyl, a substituted or
 25 unsubstituted phenyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -alkyl, phenylfuranyl or heteroaryl-

C₁-C₄-alkyl. Suitable phenyl substituents for a substituted phenyl or a substituted phenyl-C₁-C₄-alkyl include one or more of the following independently selected groups: cyano; alkyl, such as C₁-C₄-alkyl, preferably methyl; alkoxy, such as C₁-C₄-alkoxy, preferably methoxy; C₁-C₄-alkyl-S-; a halogen, preferably chloro or fluoro; a
 5 halogenated C₁-C₄-alkyl, preferably trifluoromethyl; and phenoxy. A phenoxy substituent can also be substituted with an alkyl or alkoxy group as described above. Suitable identities for R₉ include, but are not limited to, phenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, diphenylmethyl, pyrazolylmethyl, 2,4-dimethylphenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-methyl-4-methoxyphenyl, 3-
 10 methyl-4-methoxyphenyl, 4-methylthiophenyl, 3-chlorophenyl, 3-trifluoromethylphenyl, benzyl, 2-trifluoromethylbenzyl, 3-trifluoromethylbenzyl, 2-chlorobenzyl, 3-chlorobenzyl, 4-chlorobenzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-fluorobenzyl, 3-azidylphenyl, 3-(4-methoxyphenoxy)phenyl, or 5-phenylfuran-2-yl.

15 In one embodiment, R₁₀ is substituted or unsubstituted phenyl, alkyl substituted with a heteroaryl group, alkyl substituted with a heterocycloalkyl group, or an alkyl substituted with -NR₁₃R₁₄, preferably N,N-dialkylamine. Suitable identities for R₁₀ include, but are not limited to, 2-(imidazol-4-yl)ethyl, 3-(imidazol-4-yl)propyl, 3-(imidazol-1-yl)propyl, 2-(3-methylimidazol-4-yl)ethyl, 2-(morpholin-
 20 4-yl)ethyl, 2-(4-pyrazolyl)ethyl, 4-pyrazolylmethyl, 2-N,N-dimethylaminoethyl, 3-N,N-dimethylaminopropyl, and 2-(aminocarbonyl)phenyl.

R₁₁ is, preferably, a linear or branched C₁-C₄-alkyl, substituted or unsubstituted phenyl, substituted or unsubstituted benzophenonyl, pyrazolyl, aminopyrazolyl, substituted or unsubstituted indolyl-C₁-C₄-alkyl, thiophenyl,
 25 quinoxaline, substituted or unsubstituted phenyl-C₁-C₄-alkyl, pyridylcarbonylphenyl, phenylcarbonyl-C₁-C₄-alkyl, naphthyl, naphthyl-C₁-C₄-alkyl, diphenyl-C₁-C₄-alkyl, C₅-C₈-cycloalkyl-C₁-C₄-alkyl, C₁-C₄-alkylcarbonyl-C₁-C₄-alkyl, fluorenyl, pyrrolyl, N-methylpyrrolyl, or pyridyl. Suitable substituents on the phenyl ring include
 30 halogens, preferably fluoro; furanyl; thiophenyl; phenyl; benzyl; phenoxy; alkyl, such as C₁-C₄-alkyl, preferably methyl; phenoxyalkyl; C₁-C₄-alkylcarbonyl; -C(O)-benzyl; and alkoxy, such as C₁-C₄-alkoxy, preferably methoxy. Suitable substituents

-12-

on the benzophenonyl ring system include alkoxy, such as C₁-C₄-alkoxy, preferably methoxy; halogens, preferably chloro; and a C₁-C₄-alkyl group. Suitable substituents on the indole ring of a indolyl-C₁-C₄-alkyl include halogens, preferably bromo. Suitable substituents on the C₁-C₄-alkyl of a phenyl-C₁-C₄-alkyl include hydroxyl groups. Suitable substituents on the C₁-C₄-alkyl of a indolyl-C₁-C₄-alkyl include hydroxyl groups. Suitable identities of R₁₁ include, but are not limited to, benzophenon-2-yl, 4'-methoxybenzophenon-2-yl, 4'-chlorobenzophenon-2-yl, 2-(furan-2-yl)phenyl, 2-(thiophen-2-yl)phenyl, 2-benzylphenyl, 2-pyridylcarbonylphenyl, 2-(phenoxymethyl)phenyl, 2-(*t*-butylcarbonyl)phenyl, 2,2-diphenylethyl, 1-fluorenyl, (naphth-2-yl)methyl, naphth-1-yl, 3-(phenylcarbonyl)propyl, 4-phenylbutyl, 4-butylphenyl, 2-(4-chlorophenylcarbonyl)phenyl, 3-methoxyphenyl, N-methylpyrrol-2-yl, 2,3-dimethoxyphenyl, 3-butyl-2-pyridyl, 2-naphthylmethyl, 2-cyclohexylethyl, 3-methoxyphenyl, N-methyl-2-pyrrolyl, 2-cyclopentylethyl, 3-oxobutyl, 2-benzopyrazyl, quinoxalin-2-yl, 3-idolyl, (2-methylindol-3-yl)methyl, 3-(indol-3-yl)propyl, (indol-3-yl)methyl, (5-bromoindol-3-yl)methyl, 3-chlorophenyl, 3-aminopyrazol-4-yl, 2-(indol-3-yl)-1-hydroxyethyl, 3-fluorophenyl, 1-phenyl-1-hydroxymethyl, 2-phenylphenyl, 2-phenoxyphenyl, thiophen-2-yl, and isopropyl.

Certain compounds of formula I, II or III may contain one or more chiral centres, and exist in different optically active forms. When compounds of formula I contain one chiral centre, the compounds exist in two enantiomeric forms and the present invention includes both enantiomers and mixtures of enantiomers. The enantiomers may be resolved by methods known to those skilled in the art, for example by formation of diastereoisomeric salts which may be separated, for example, by crystallization; formation of diastereoisomeric derivatives or complexes which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic esterification; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, for example silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described

above, a further step is required to liberate the desired enantiomeric form. Alternatively, specific enantiomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer into the other by asymmetric transformation.

- 5 When a compound of formula I, II or III contains more than one chiral centre it may exist in diastereoisomeric forms. The diastereoisomeric pairs may be separated by methods known to those skilled in the art, for example chromatography or crystallization and the individual enantiomers within each pair may be separated as described above. The present invention includes each diastereoisomer of compounds of formula I, II or III
- 10 and mixtures where the diastereoisomers are unresolved.

Certain compounds of formula I, II, or III may exist in different tautomeric forms or as different geometric isomers, and the present invention includes each tautomer and/or geometric isomer of compounds of formula I, II or III and tautomeric mixtures thereof.

- 15 Certain compounds of formula I, II or III may exist in different stable conformational forms which may be separable. Torsional asymmetry due to restricted rotation about an asymmetric single bond, for example because of steric hindrance or ring strain, may permit separation of different conformers. The present invention includes each conformational isomer of compounds of formula I, II or III and mixtures
- 20 of conformational isomers thereof.

Certain compounds of formula I, II or III may exist in zwitterionic form and the present invention includes each zwitterionic form of compounds of formula I, II or III and mixtures thereof.

- 25 The present invention further relates to pharmaceutically acceptable salts of the compounds of Formulas I, II and III. The phrase a "pharmaceutically acceptable salt" is intended to include a salt which retains the biological effectiveness and properties of the free base and which can be obtained by reaction with an inorganic or organic acid, such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, organic sulfonic acid, organic carboxylic acid, organic
- 30 phosphoric acid, for example, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, lactic acid, tartaric acid and the like.

Compounds of Formulas I, II and III are modulators of the signaling processes which follow binding of TNF- α to its receptor. In preferred embodiments, the compounds of the invention interrupt the signaling process. Without being bound by theory, it is believed that these compounds interfere with one or more steps of the signaling cascade which includes TNF- α . Thus, these compounds are effective as therapeutic agents for medical conditions in which one or more signaling processes involving TNF- α play a role and include, for example, conditions in which excessive TNF- α is produced. When selecting substituents for the various compounds of Formulas I, II and III, the ordinarily skilled artisan can utilize available information to ensure selection of a stable compound with substituents that will enhance the desired therapeutic activity. The present compounds can be administered to a patient having a TNF- α mediated medical condition, for example, to inhibit the development of the condition, to inhibit its further progression, and/or to ameliorate the symptoms associated with the condition.

Thus, in one embodiment, the present invention relates to a method of treating a TNF- α mediated condition in a patient. The method comprises the step of administering to the patient a therapeutically effective amount of at least one compound of Formula I, Formula II or Formula III, as described above. The patient can be a human or any animal which is suffering from a TNF- α mediated condition or which is believed to be susceptible to development of a TNF- α mediated condition. Preferably, the patient is a domestic animal, such as a fowl, for example, a chicken, or a mammal, for example, a bovine, porcine, canine, feline or equine animal. More preferably, the patient is a human.

The language a "TNF- α mediated condition" is intended to include a medical condition, such as a chronic or acute disease or pathology, or other undesirable physical state, in which a signaling cascade including TNF- α plays a role, whether, for example, in development, progression or maintenance of the condition. Examples of TNF- α mediated conditions include, but are not limited to:

(A) acute and chronic immune and autoimmune pathologies, such as systemic lupus erythematosus (SLE), rheumatoid arthritis, thyroidosis, graft versus host disease,

-15-

scleroderma, diabetes mellitus, Graves' disease, and the like;

(B) infections, including sepsis syndrome, circulatory collapse and shock resulting from acute or chronic bacterial infection, acute and chronic parasitic infection, and/or infectious diseases, whether bacterial, viral or fungal in origin, such as a
5 HIV or AIDS, and including symptoms of cachexia, autoimmune disorders, Acquired Immune Deficiency Syndrome, dementia complex and infections;

(C) inflammatory diseases, such as chronic inflammatory pathologies, including sarcoidosis, chronic inflammatory bowel disease, ulcerative colitis and Crohn's pathology, and vascular inflammatory pathologies, such as, disseminated
10 intravascular coagulation, atherosclerosis and Kawasaki's pathology;

(D) neurodegenerative diseases, including, demyelinating diseases, such as multiple sclerosis and acute transverse myelitis; extrapyramidal and cerebellar disorders such as lesions of the corticospinal system; disorders of the basal ganglia or cerebellar disorders; hyperkinetic movement disorders such as Huntington's Chorea and senile
15 chorea; drug-induced movement disorders, such as those induced by drugs which block CNS dopamine receptors; hypokinetic movement disorders, such as Parkinson's disease; progressive supranuclear palsy; Cerebellar and Spinocerebellar Disorders, such as structural lesions of the cerebellum; spinocerebellar degenerations, such as spinal ataxia, Friedreich's ataxia, cerebellar cortical
20 degenerations; multiple systems degenerations, such as Mancel, Dejerine-Thomas, Shi-Drager, and Machado-Joseph; systemic disorders, such as Refsum's disease, abetalipoproteinemia, ataxia, telangiectasia, and mitochondrial multisystem disorder; demyelinating core disorders, such as multiple sclerosis, acute transverse myelitis; disorders of the motor unit, such as neurogenic muscular atrophies, such as anterior
25 horn cell degeneration, amyotrophic lateral sclerosis, infantile spinal muscular atrophy and juvenile spinal muscular atrophy; Alzheimer's disease; Down's Syndrome in middle age; Diffuse Lewy body disease; Senile Dementia of Lewy body type; Wernicke-Korsakoff syndrome; chronic alcoholism; Creutzfeldt-Jakob

-16-

disease; Subacute sclerosing panencephalitis, Hallerorden-Spatz disease; and Dementia pugilistica, or any subset of conditions, symptoms or syndromes thereof;

(E) malignant pathologies involving TNF- α secreting tumors or other malignancies involving TNF, such as leukemias including acute, chronic myelocytic, chronic lymphocytic and/or myelodysplastic syndrome; lymphomas including Hodgkin's and non-Hodgkin's lymphomas; and malignant lymphomas, such as Burkitt's lymphoma or Mycosis fungoides; and

(F) alcohol-induced hepatitis.

See, e.g., Berkow, *et al.*, eds., *The Merck Manual*, 16th edition, chapter 11, pp 1380-1529, Merck and Co., Rahway, N.J., (1992), which reference, and references cited therein, are entirely incorporated herein by reference.

The language a "therapeutically effective amount" is intended to include an amount which is sufficient to inhibit, totally or partially, the TNF- α mediated condition, prevent its further progression or ameliorate the symptoms associated with the condition. Such an amount, when administered prophylactically to a patient thought to be susceptible to development of a TNF- α mediated condition, can also be effective to prevent or lessen the severity of the TNF- α mediated condition.

The compounds of this invention can be administered to the patient by themselves or in pharmaceutical compositions where they are mixed with a suitable carrier or excipient at doses to treat or ameliorate the symptoms associated with the TNF- α mediated condition. Mixtures of these compounds can also be administered to the patient as a simple mixture or in suitable formulated pharmaceutical compositions. Techniques for formulation and administration of the compounds of the instant application can be found in *Remington: the Science and Practice of Pharmacy*, 19th edition, Mack Publishing Co., Easton, PA. (1995).

Suitable routes of administration can, for example, include oral, eyedrop, rectal, transmucosal, topical, or intestinal administration; parenteral delivery,

including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

The compounds of the invention can also be administered in a targeted drug delivery system, such as, for example, in a liposome coated with endothelial cell-specific antibody.

The invention also relates to the use of a compound of Formula I, Formula II or Formula III, as described above, for the manufacture of a medicament for treating a TNF- α mediated condition.

The pharmaceutical compositions of the present invention can be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention thus can be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

Pharmaceutical preparations for oral use can be obtained by combining the active compound with a solid excipient, optionally grinding a resulting mixture, and

-18-

processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol, cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in a conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of a dry powder inhaler, or an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of pressurized aerosol the dosage unit may be determined by providing a valve to

deliver a metered amount. Capsules and cartridges of gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds can be formulated for parenteral administration by injection, including bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, such as in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

10 Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes.

15 Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

20 Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, such as sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

25 In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation, for example, subcutaneously or intramuscularly or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials, for example, as an emulsion in an acceptable oil, or ion exchange resins, or as sparingly soluble derivatives, for
30 example, as a sparingly soluble salt.

The pharmaceutical compositions can also include suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

5 Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. More specifically, a therapeutically effective amount, as previously defined, denotes an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the
10 effective amounts is well within the capability of those skilled in the relevant art.

For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from *in vitro* assays and animal models. For example, a dose can be formulated in cellular and animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cellular
15 assays, *i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of TNF- α activity. In some cases it is appropriate to determine the IC_{50} in the presence of 3 to 5% serum albumin, since such a determination approximates the binding effects of plasma protein on the compound. Such information can be used to more accurately determine useful doses in humans. Further, the most preferred
20 compounds for systemic administration effectively inhibit TNF- α signaling in intact cells at levels that are safely achievable in plasma.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the maximum tolerated dose (MTD) and the ED_{50} (effective dose for
25 50% maximal response). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between MTD and ED_{50} . Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within
30 a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed

and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See, *e.g.*, Fingl, *et al.*, 1975, in *The Pharmacological Basis of Therapeutics*, Chapter 1, p. 1). In the treatment of crises, the administration of an acute bolus or an infusion approaching the MTD may be required to obtain a rapid response.

Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the desired effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data, *e.g.*, the concentration necessary to achieve 50-90% inhibition of protein kinase using the assays described herein. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.


Dosage intervals can also be determined using the MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90% until the desired amelioration of symptoms is achieved. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

The amount of composition administered will be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

SYNTHETIC STRATEGIES

In general, the substituted pyrazole ring of Formula I can be prepared by reaction of a 1,3-dicarbonylalkane with a hydrazine (Scheme I).

-22-




Scheme I: Formation of pyrazole ring.

When the method in Scheme I is used to construct the compounds represented by Formula I, R₂₈ is ring A and R₂₇ is -X(CH₂)_n-Y. R₂₅ and R₂₆ are each, independently,
 5 -H or a substituted or unsubstituted alkyl or a substituted or unsubstituted alkoxy.
 The above reaction was used to prepare the compounds in Examples 6-10 and 12-15.

The reaction to form the pyrazole ring is carried out in a polar solvent, such as water, an alcohol or an ether. Preferably, the reaction is carried out in an alcohol, such as ethanol. The reaction temperature is about 35°C to about 150°C, preferably
 10 about 70°C to about 90°C. Typically, the reaction is carried out at the reflux temperature of the solvent used.

In compounds which can be represented by Formula I, ring A can be a pyridine, a pyrimidine or a triazine ring. Substituted 2-hydrazinopyrimidines can be prepared by addition of thiourea to 3-(N, N-dimethylamino)prop-2-en-1-one in the
 15 presence of a base, followed by *in situ* methylation of the cyclic thiourea formed (see Scheme II, step 1). The resultant 4-substituted 2-methylsulfanylpurimidine is treated with hydrazine and heat to form a 4-substituted 2-hydrazinopyrimidines (see Scheme II, step 2).

-23-



Scheme II: Preparation of substituted 2-hydrazinopyrimidine.

The reaction depicted in Scheme II was used in Example 5 to prepare the intermediate, 2-hydrazino-4-(pyridin-2-yl)-pyrimidine. This intermediate was reacted with a 1,3-dicarbonylalkane via the method shown in Scheme I to prepare a substituted pyrazole in Examples 6-9.

The reaction to form substituted 2-methylsulfanylpurine (see Scheme II, step 1) is carried out in a polar solvent, such as water, an alcohol or an ether. Preferably, the reaction is carried out in an alcohol, such as ethanol, using an alkaline or alkaline earth metal alkoxide as the base. The reaction temperature is about 35°C to about 150°C, preferably about 70°C to about 90°C. Typically, the reaction is carried out at the reflux temperature of the solvent used. The reaction is usually allowed to proceed for about 1 hour to about 24 hours, preferably about 2 to


about 5 hours at an elevated temperature, then the reaction is cooled and a primary haloalkane, preferably an iodoalkane, is added to alkylate the thiol group.

The substituted 2-methylsulfanylpurine is then treated with hydrazine at elevated temperatures to form substituted 2-hydrazinopurine. The reaction can
5 be carried out in a polar solvent, such as water, an alcohol or an ether. Typically, the reaction is carried out in water having about 35% (vol/vol) hydrazine. The reaction temperature is about 80°C to about 110°C, preferably, about 100°C.

Substituted 2-hydrazinotriazines can be prepared by reacting 2-
amidopyridine or 2-amidoquinoxaline with an alkoxy-*bis*-(dimethylamino)methane,
10 such as *tert*-butoxy-*bis*-(dimethylamino)methane, at a temperature of about 80°C to about 170°C, preferably at about 145°C to about 160°C for about 1 hour to about 5 hours, preferably about 2 hours to about 4 hours to yield a first intermediate. The reaction is typically carried out in a polar aprotic solvent such as dimethylformamide. The 2-amidopyridine or 2-amidoquinoxaline is typically
15 present in a concentration of about 0.2 M to about 1 M and the alkoxy-*bis*-(dimethylamino)methane is present in about 1.5 equivalents to about 3 equivalents in relation to the 2-amidopyridine or 2-amidoquinoxaline (Scheme III, step 1a).

The first intermediate is then contacted with thiourea and a base in a polar solvent such as water, an alcohol or an ether. Preferably, the reaction is carried out
20 in an alcohol, such as ethanol, using an alkaline or alkaline earth metal alkoxide as the base. The reaction temperature is about 35°C to about 150°C, preferably about 70°C to about 90°C. The reaction is usually allowed to proceed for about 1 hour to about 24 hours, preferably about 2 to about 5 hours at an elevated temperature to form a 4-substituted-2-thiotriazine (see Scheme III, step 1b). The 4-substituted-2-
25 thiotriazine is then alkylated and substituted with hydrazine (see Scheme III, steps 2 and 3) to form a substituted 2-hydrazinotriazine via the method described above for forming substituted 2-hydrazinopurine.

-25-




Scheme III: Formation of a substituted 2-hydrazinotriazine.

2-Hydrazinotriazine can react with a 1,3-dicarbonylalkane to form a pyrazole ring as shown in the method of Scheme I. Examples 12-14 were prepared via the method of Schemes III followed by the method of Scheme I.

When ring A is a pyridine ring, the pyrazole ring can be formed first by reaction of hydrazine with 1,3-dicarbonylalkane via the method shown in Scheme I where R₂₈ is a hydrogen. The pyrazole ring can then be added to the pyridine ring via a substitution reaction (see Scheme IV).

-26-



Scheme IV: Formation of a substitute 1-(pyridin-2-yl)pyrazole.


In Scheme IV, R₂₉ is a leaving group, such as a halogen. The pyrazole is contacted with a base that is sufficiently strong to deprotonate the pyrazole for about 3 minutes to about 30 minutes prior to addition of the substituted pyridine. Typically, a
 5 hydride salt, such as sodium hydride is used. After addition of the substituted pyridine the reaction is heated for about 6 hours to about 24 hours at a temperature of about 80°C to about 120°C to form a 6-substituted 2-(pyrazol-1-yl)pyridine. The reaction is typically carried out in an aprotic solvent such as an ether. Example 10

-27-

was prepared using the method of Scheme I followed by the method of Scheme IV.

The compounds listed in Tables 2 and 4 were prepared using the methods of Schemes I-IV.

The substituted triazole ring of Formula II can be prepared by reaction of a
5 N-thiocarbonylamide with a hydrazine (see Scheme V).




Scheme V: Formation of a triazole.

When the reaction in Scheme V is used to form the compounds of Formula II, R_{24} is a 4-(pyridin-2-yl)triazin-2-yl. The reaction is carried out in an acidic buffer, such as
10 carboxylic acid and the salt of the carboxylic acid, for example acetic acid and sodium acetate. An organic solvent can also be present, such as an ether. The reaction is heated to about 70°C to about 110°C for about 6 hours to about 24 hours. Example 11 is representative of the method depicted in Scheme V.

The compounds in Table 3 were prepared using the methods of Schemes III
15 and V.

The compounds represented by Formula III can be formed via an Ugi reaction, as shown in Scheme VI.

-28-




Scheme VI: Ugi reaction to form the compounds represented by Formula III.

For reviews of Ugi reactions see Gross and Meienhofer, *The Peptides*, vol. 2, pp. 365-381, Academic Press, New York, (1980); *Intra-Sci. Chem. Rep.* (1971), 5:229-261; *Rec. Chem. Prog.* (1969), 30:289-311; and Eberle, *et al.*, *Tetrahedron* (1978), 34:977, the entire teachings of which are incorporated herein by reference. The starting materials for the Ugi reaction, *i.e.*, an isonitrile, a carboxylic acid, an aldehyde or a ketone and an amine, are mixed together in a polar solvent such as an alcohol, preferably methanol or ethanol. The reaction is heated to about 40°C to about 80°C for about 6 hours to about 24 hours. Examples 1 and 2 are representative of compounds prepared using an Ugi reaction.

A limitation of the Ugi reaction is that the terminal nitrogen atom, *i.e.*, the nitrogen substituted with R₈ is monosubstituted. Therefore, a second method of forming the compounds represented by Formula III was developed wherein the terminal nitrogen can be mono- or disubstituted (see Scheme VII).

-29-



Scheme VII: Formation of compounds represented by Formula III.


In Scheme VII, R_{29} is a leaving group, such as a halide, and R_{30} is a substituted or unsubstituted aryl or a substituted or unsubstituted alkyl. The starting materials for the first step are mixed together in about equal molar amounts in a nonpolar, aprotic solvent such as methylene chloride, or an ether, for about 0.5 hours to about 6 hours, preferably about 1 hour to form an acetoxamide. This is followed by hydrolysis with lithium hydroxide to form an α -hydroxyamide.

In step 2, the α -hydroxyamide formed in step 1 is dissolved in an aprotic solvent, such as an ether, at a concentration of about 0.2 M to about 0.4 M and is treated with about 1.1 equivalents to about 1.5 equivalents, preferably about 1.2 equivalents, of a strong base, such as a hydride salt, for example, sodium hydride, 5 for about 2 minutes to about 20 minutes, followed by addition of about 1.1 equivalents to about 1.5 equivalents, preferably about 1.2 equivalents, of an arylsulfonyl halide or alkylsulfonyl halide. After addition of the arylsulfonyl halide or alkylsulfonyl halide, the reaction is allowed to proceed for about 1 hour to about 6 hours, then about 3 equivalents to about 5 equivalents, preferably about 4 10 equivalents, of a primary amine is added to the reaction mixture accompanied by addition of a polar solvent such as an alcohol. After addition of the primary amine, the reaction is allowed to proceed for about 6 hours to about 24 hours.

The product of step 2 is treated with about 1.1 equivalents to about 1.8 equivalents of an acid halide in a non-polar solvent, such as a halogenated alkane or 15 an ether in the presence of a base, such as a tertiary amine, for about 1 hour to about 4 hours at a temperature of about 30°C to about 80°C to form the desired product. Example 4 is representative of this method.

A third method of forming the compounds represented by Formula III was developed wherein the stereochemistry of the carbon substituted with R₉ can be 20 controlled (see Scheme VIII).

-31-



Scheme VIII: Third method of forming the compounds represented by Formula III.

- In Scheme VIII, R_{29} is a leaving group, such as a halogen and R_{31} is an amine protecting group. Methods for protecting amines can be found in Greene, *et al.*, *Protecting Groups in Organic Synthesis*, 2nd Edition, John Wiley & Sons, Inc., (1991), pp. 309-405, the entire teachings of which are incorporated herein by reference. The starting material for the synthesis depicted in Scheme VIII can be a natural or unnatural amino acid wherein the amine group is protected. Preferred

protecting groups for the amine are 9-fluorenylmethyl carbamate and *t*-butyl carbamate. When a chiral amino acid is the starting material, the chiral center at the α -carbon is retained through out the synthesis.

In step 1 of Scheme VIII, the carboxylic acid of the starting material is transformed into an activated ester by methods known to those skilled in the art. The activated ester is then contacted with a primary amine to form an amide. In step 2, a Mitsunobu reaction is used to displace a hydroxyl group of R_{10} -OH with the protected amine of the product formed in step 1 by treating the alcohol with triphenylphosphine and di-*tert*-butyl azodicarboxylate in the presence of the product of step 1. Fukuyama T., *et al.*, *Tetrahedron Letters* (1995), 36:6373. The amine is then deprotected in step 3 and reacted with an acid halide as described above for step 3 of synthetic Scheme VII. Example 3 is representative of this method.

The compounds of Table I were formed using the methods of Schemes VI-VIII.

15 EXAMPLES

I. SYNTHETIC METHODS

Example 1: 2-Benzoyl-*N*-[(2,2-diphenyl-ethylcarbamoyl)-phenyl-methyl]-*N*-[2-(1*H*-imidazol-4-yl)-ethyl]-benzamide (Compound 5)

Step 1) Preparation of 2,2-Diphenylethylisocyanide.

20 A stirred suspension of 2,2-diphenylethylamine (7.00 g, 35.5 mmol) in ethyl formate (25 mL) was heated at reflux overnight. The reaction was concentrated to afford the corresponding formamide as an off-white solid. The crude product was taken up in methylene chloride (50 mL) and diisopropylamine (13.3 mL, 94.9 mmol) and cooled in an ice bath. While stirring, phosphorus oxychloride (5.0 mL, 54 mmol) was added. One hour later, the ice bath was removed and aqueous sodium carbonate solution (75 mL), water (50 mL) and methylene chloride (50 mL) were added. The mixture was stirred vigorously for 1 hour. The organic layer was

-33-

washed with water, dried (sodium sulfate) and concentrated. The resulting crude product was filtered through a plug of silica (methylene chloride) to afford 6.62 g (90%) of product as an off-white solid: ^1H NMR (CDCl_3) δ 7.38-7.18 (m, 10H), 4.35 (t, $J = 7.5$ Hz, 1H), 3.98 (d, $J = 7.3$, 2H) ppm.

5 Step 2) Preparation of Compound 5.

A solution of histamine (1.11 g, 10.0 mmol), 2-benzoylbenzoic acid (2.26 g, 10.0 mmol), benzaldehyde (1.06 g, 10.0 mmol) and the product of step 1 (2.07 g, 10.0 mmol) in methanol was heated at reflux overnight. The reaction was concentrated and the residue partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was dried (sodium sulfate) and concentrated to afford a foamy light brown solid. Flash chromatography over silica (methylene chloride/methanol) afforded 2.99 g (47%) of the product as a beige solid: ^1H NMR (CDCl_3) δ 7.88-6.84 (m, 25H), 6.65-6.25 (m, 2H), 5.66-5.30 (m, 1H), 4.44-4.20 (m, 1H), 4.17-3.84 (m, 2H), 3.47-3.03 (m, 2H), 2.69-2.18 (m, 2H) ppm. MS (ESI) m/z 634 ($\text{M}+\text{H}^+$).

Example 2: *N*-[(4-Cyano-phenyl)-(2,2-diphenyl-ethylcarbamoyl)-methyl]-2-(2,2-dimethyl-propionyl)-*N*-[2-(1*H*-imidazol-4-yl)-ethyl]-benzamide
(Compound 54)

A solution of histamine (0.100 g, 0.900 mmol), 2-pivaloylbenzoic acid (0.186 g, 0.902 mmol), 3-cyanobenzaldehyde (0.118 g, 0.900 mmol) and the product of step 1 of example 1 (0.187 g, 0.902 mmol) in methanol was heated at reflux overnight. The reaction was concentrated and the residue partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was dried (sodium sulfate) and concentrated to afford 0.521 g (91%) of crude product as a foamy amber solid. This material was used for biological testing without further purification.

Example 3: 2-Benzoyl-*N*-[(1'*R*)-1'-(2',2'-diphenyl-ethylcarbamoyl)-2'-phenyl-ethyl]-*N*-[2-(1*H*-imidazol-4-yl)-ethyl]-benzamide (Compound 47)

Step 1) Preparation of (2*R*)-2-Amino-*N*-(2,2-diphenyl-ethyl)-3-phenyl-propionamide.

5 To a stirred solution of *N*-(*tert*-butoxycarbonyl)-*D*-phenylalanine (11.0 g, 41.5 mmol) in chloroform (125 mL) was added 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (7.95 g, 41.5 mmol) and 1-hydroxybenzotriazole (5.60 g, 41.4 mmol). After 5 minutes, 2,2-diphenylethylamine (7.44 g, 37.7 mmol) was added. The reaction was stirred for 2 hours, diluted with chloroform (100 mL)
10 and washed with aqueous sodium bicarbonate solution. The organic layer was dried (sodium sulfate) and concentrated to afford an off-white solid. Trituration with ether afforded the crude amide. This material was taken up in methylene chloride (100 mL) and treated with trifluoroacetic acid (45 mL, 580 mmol). After stirring for 15 minutes, the reaction was concentrated and the residue was partitioned between ethyl
15 acetate and aqueous sodium bicarbonate. The organic layer was dried (sodium sulfate) and concentrated to afford 12.4 g (111% yield inflated by entrained residual solvent) of product as a colorless gum: ¹H NMR (CDCl₃) δ 7.34-6.95 (m, 15H), 4.10 (t, *J* = 8.2, 1H), 3.88-3.79 (m, 2H), 3.68-3.61 (m, 1H), 3.49 (br s, 2H), 3.13-3.03 (m, 1H), 2.70-2.59 (m, 1H) ppm.

20 Step 2) Preparation of (2*R*)-*N*-(2,2-Diphenyl-ethyl)-2-(2-nitrobenzenesulfonylamino)-3-phenyl-propionamide.

To a stirred solution of the product of step 1 (12.3 g, 35.6 mmol) in methylene chloride (200 mL) was added 2-nitrobenzenesulfonyl chloride (9.00 g, 40.6 mmol) followed by triethylamine (6.4 mL, 46 mmol). After 30 minutes the
25 reaction was concentrated and the residue partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was dried (sodium sulfate)

and concentrated. The resulting amber foam was purified by flash chromatography over silica (hexane/ethyl acetate) to afford 14.0 g (74%) of product as a colorless foam: ¹H NMR (CDCl₃) δ 7.95-7.87 (m, 1H), 7.76-7.62 (m, 3H), 7.35-7.15 (m, 10H), 7.06-6.94 (m, 3H), 6.90-6.83 (m, 2H), 6.42 (t, *J* = 5.6, 1H), 5.80 (d, *J* = 5.8, 1H), 4.12 (t, *J* = 8.1, 1H), 4.01-3.82 (m, 3H), 3.18-3.08 (m, 1H), 2.68-3.57 (m, 1H) ppm.

Step 3) Preparation of (2*R*)-*N*-(2,2-Diphenyl-ethyl)-2-[(2-nitro-benzenesulfonyl)-[2-(1-trityl-1*H*-imidazol-4-yl)-ethyl]-amino}-3-phenyl-propionamide.

To a stirred solution of the product of step 2 (3.40 g, 6.42 mmol) and 2-(1-trityl-1*H*-imidazol-4-yl)-ethanol (2.73 g, 7.70 mmol) in methylene chloride (40 mL) was added triphenylphosphine (2.19 g, 8.35 mmol) followed by di-*tert*-butyl azodicarboxylate (1.77 g, 7.69 mmol). After 3 hours the reaction was concentrated and the residue partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was dried (sodium sulfate) and concentrated to yield an amber gum. Flash chromatography over silica (methylene chloride/methanol) afforded 5.42 g (97%) of partially purified product (contaminated with di-*tert*-butyl hydrazodiformate) as a pale yellow foam: ¹H NMR (CDCl₃) δ 7.81-7.74 (m, 1H), 7.64-7.55 (m, 1H), 7.53-7.44 (m, 2H), 7.41-7.03 (m, 31H), 6.69 (t, *J* = 5.6, 1H), 6.53 (s, 1H), 4.48 (t, *J* = 8.1, 1H), 4.08 (t, *J* = 8.1, 1H), 3.84-3.58 (m, 4H), 3.44-3.34 (m, 1H), 2.91-3.80 (m, 1H), 2.78-2.68 (m, 1H), 2.63-2.51 (m, 1H) ppm.

Step 4) Preparation of (2*R*)-*N*-(2,2-Diphenyl-ethyl)-3-phenyl-2-[2-(1-trityl-1*H*-imidazol-4-yl)-ethylamino]-propionamide.

To a stirred solution of the product of step 3 (5.42 g, 6.26 mmol) in *N,N*-dimethylformamide (17 mL) was added mercaptoacetic acid (1.1 mL, 15.8 mmol) followed by lithium hydroxide hydrate (1.31 g, 31.2 mmol). The mixture was stirred for 3 hours and then partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was combined with a second ethyl acetate

-36-

extract, dried (sodium sulfate) and concentrated to furnish an amber oil. Flash chromatography over silica (methylene chloride/methanolic ammonia solution) provided 2.50 g (59%) of product as a colorless tacky foam: ¹H NMR (CDCl₃) δ 7.54 (t, *J* = 5.5, 1H), 7.38-7.01 (m, 31H), 6.33 (s, 1H), 4.16 (t, *J* = 8.3, 1H), 4.02-
5 3.92 (m, 1H), 3.83-3.74 (m, 1H), 3.22-3.15 (m, 1H), 3.12-3.04 (m, 1H), 2.51-2.18 (m, 5H) ppm.

Step 5) Preparation of 2-Benzoyl-*N*-[(1'*R*)-1-(2',2'-diphenyl-ethylcarbamoyl)-2'-phenyl-ethyl]-*N*-[2-(1-trityl-1*H*-imidazol-4-yl)-ethyl]-benzamide.

2-Benzoylbenzoic acid (1.33 g, 5.88 mmol) was taken up in a 2M solution of
10 oxalyl chloride in methylene chloride (4.5 mL, 9.0 mmol). One drop of *N,N*-dimethylformamide was added and the frothy mixture was stirred for 2 hours before concentrating. To the residue was added a solution of the product of step 4 (2.50 g, 3.67 mmol) in chloroform (30 mL) and triethylamine (1.05 mL, 7.53 mmol). The mixture was heated at reflux for 2.5 hours. At this time, more triethylamine (0.50
15 mL, 3.6 mmol) was added and the reaction was heated for additional 30 minutes. The reaction was then concentrated and the residue was partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was dried (sodium sulfate) and concentrated to afford an amber foam. This material was filtered through silica (methylene chloride/methanolic ammonia solution) to afford
20 1.44 g (44%) of a pale amber foam which was used without further purification in the next step.

Step 6) Preparation of Compound 47.

A stirred solution of the product of step 5 (1.44 g, 1.62 mmol) in 4:1 acetic acid/water (7 mL) was heated in a water bath (90°C) for 30 minutes. The reaction
25 solution was then cooled and partitioned between ethyl acetate and aqueous sodium carbonate solution. The organic layer was combined with a second ethyl acetate extract, dried (sodium sulfate) and concentrated to afford an amber foam. Flash

-37-

chromatography over silica (methylene chloride/methanol) afforded 0.62 g (59%) of product as a foamy pale amber solid. Comparisons by thin layer chromatography and ¹H NMR spectroscopy indicated that this material is identical to the corresponding racemic Ugi product, compound 43. Analytical chiral HPLC (ChiralPack AD, 4.6 x 250 mm; eluant: 0.1% diethylamine in 88:12 hexane/ethanol; flow: 1 mL/min; detection: 260 nM) indicated that the material is a single enantiomer within detection limits: ¹H NMR (CDCl₃) δ 8.35 (br s, 1H), 7.90-6.94 (m, 27H), 6.78-6.57 (m, 1H), 6.39-6.17 (m, 1H), 4.65-3.72 (m, 4H), 3.64-2.70 (m, 5H), 2.46-2.18 (m, 1H) ppm. MS (ESI) *m/z* 648 (M+H⁺).

10 Example 4: 2-Benzoyl-*N*-{[(2,2-diphenyl-ethyl)-methyl-carbamoyl]-phenyl-methyl}-*N*-[2-(1*H*-imidazol-4-yl)-ethyl]-benzamide (Compound 26)

Step 1) Preparation *N*-(2,2-Diphenyl-ethyl)-2-hydroxy-*N*-methyl-2-phenyl-acetamide.

To a stirred solution of (2,2-diphenylethyl)methylamine (4.97 g, 23.5 mmol) in chloroform (50 mL) was added O-acetylmandelic chloride (5.3 mL, 24 mmol) followed by triethylamine (4.0 mL, 29 mmol). The reaction was stirred for 1 hour and then concentrated. The residue was partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was dried (sodium sulfate) and concentrated to afford a colorless gum. The crude amide was taken up in 1:1 tetrahydrofuran/methanol (80 mL) and treated with 1 N aqueous lithium hydroxide (30 mL, 30 mmol). The mixture was stirred for 2 hours and concentrated to remove the organic solvents. The remaining aqueous solution was diluted with water and extracted twice with ethyl acetate and once with methylene chloride. The combined organic extracts were dried (sodium sulfate) and concentrated to afford an off-white solid. This material was triturated with diethyl ether to afford 7.59 g (93%) of product as a white solid: ¹H NMR (CDCl₃) δ 7.43-6.98 (m, 15H), 5.07 (s, 1H), 4.57-4.44 (m, 1H), 4.36-4.28 (m, 1H), 3.76-3.64 (m, 1H), 2.46 (s, 3H) ppm.

Step 2) Preparation of *N*-(2,2-Diphenyl-ethyl)-2-[2-(1*H*-imidazol-4-yl)-ethylamino]-*N*-methyl-2-phenyl-acetamide.

To stirred solution of the product of step 1 (1.50 g, 4.34 mmol) in tetrahydrofuran (15 mL) was added a 60% dispersion of sodium hydride in mineral
5 oil (0.210 g, 5.25 mmol). After gas evolution ceased, *p*-toluenesulfonyl chloride (0.990 g, 5.20 mmol) was added and the mixture was stirred for 2 hours. At this time, histamine (1.93 g, 17.4 mmol) was added followed by methanol (10 mL). The reaction was stirred overnight and concentrated. The residue was partitioned between ethyl acetate and aqueous sodium carbonate solution. The organic layer
10 was dried (sodium sulfate) and concentrated to afford a pasty white solid. Flash chromatography over silica (methylene chloride/methanolic ammonia solution) provided 1.08 g (57%) of product as a colorless tacky foam: ¹H NMR (CDCl₃) δ 7.52-7.43 (m, 1H), 7.40-7.00 (m, 15H), 6.78-6.69 (m, 1H), 4.36-3.97 (m, 4H), 2.92-2.20 (m, 7H) ppm.

15 Step 3) Preparation of Compound 26.

2-Benzoylbenzoic acid (0.341 g, 1.51 mmol) was taken up in a 2M solution of oxalyl chloride in methylene chloride (1.5 mL, 3.0 mmol). One drop of *N,N*-dimethylformamide was added and the frothy mixture was stirred overnight and concentrated. A solution of the crude 2-benzoylbenzoyl chloride in methylene
20 chloride (4 mL) followed by triethylamine (0.25 mL, 1.8 mmol) was added to the product of step 2 (0.220 g, 0.502 mmol). The mixture was heated at reflux for 2 hours and concentrated. The residue was taken up in 5:1 methanol/1 N aqueous lithium hydroxide (12 mL) and stirred for 1 hour. The reaction was again concentrated and the residue was partitioned between ethyl acetate and aqueous
25 sodium bicarbonate solution. The organic layer was dried (sodium sulfate) and concentrated to afford a foamy amber solid. Flash chromatography over silica (methylene chloride/methanol) provided 0.229 g (71%) of product as a foamy pale

amber solid: ^1H NMR (CDCl_3) δ 7.91-6.92 (m, 26H), 6.54-6.39 (m, 1H), 6.34-6.06 (m, 1H), 4.83-4.31 (m, 2H), 3.97-2.82 (m, 3H), 2.70-2.39 (m, 3H), 2.35-2.09 (m, 2H) ppm. MS (ESI) m/z 648 ($\text{M}+\text{H}^+$).

Example 5: 2-Hydrazino-4-(Pyridin-2-yl)-Pyrimidine

5

Step 1) Preparation of 2-Methylsulfanyl-4-pyridin-2-yl-pyrimidine.

(Synthetic intermediate used in Examples 6, 7, 8 & 15)

To a stirred solution of sodium ethoxide (freshly prepared from sodium metal (5.11 g, 222 mmol) and ethanol (350 mL)) was added 3-dimethylamino-1-pyridin-2-yl-propenone (28.0 g, 159 mmol) and thiourea (12.9 g, 170 mmol). The solution
10 was heated to reflux and stirred for 3 hours. The solution was allowed to cool to room temperature, and then iodomethane (13.8 mL, 222 mmol) was added dropwise over 10 minutes. The solution was stirred at room temperature for 2 hours, and then diluted with saturated ammonium chloride solution (250 mL) and water (250 mL).
15 The suspension was extracted with ethyl ether (200 mL x 3), and the combined organic extracts washed with saturated sodium thiosulfate solution (150 mL) and brine (150 mL). The organic phase was dried (magnesium sulfate), filtered, and concentrated to provide 30.1 g (96%) of the product as a brown oil: ^1H NMR (CDCl_3) δ 8.70-8.69 (m, 1H), 8.64 (d, $J = 5.2$ Hz, 1H), 8.48 (d, $J = 7.9$ Hz, 1H),
20 8.01 (d, $J = 5.2$ Hz, 1H), 7.87-7.82 (m, 1H), 7.41-7.38 (m, 1H), 2.65 (s, 3H) ppm. ^{13}C NMR (CDCl_3) δ 172.6, 163.1, 158.5, 153.9, 149.7, 137.3, 125.7, 122.0, 112.6, 14.5 ppm.

Step 2) Preparation of (4-Pyridin-2-yl-pyrimidin-2-yl)-hydrazine.

To hydrazine hydrate (90 mL) was added 2-methylsulfanyl-4-pyridin-2-yl-pyrimidine (30.0g, 148 mmol). The solution was heated to reflux and stirred for 17
25 hours. The solution was allowed to cool to room temperature, and a yellow precipitate formed. The solids were removed by filtration, washed with water, and

dried to provide 22.5 g (81%) of the product as yellow micro needles: ^1H NMR (DMSO- d_6) δ 8.68-8.67 (m, 1H), 8.45-8.41 (m, 2H), 8.29 (s, 1H), 7.98-7.93 (m, 1H), 7.52-7.48 (m, 2H), 4.28 (s, 2H) ppm. ^{13}C NMR (DMSO- d_6) δ 164.6, 162.6, 159.3, 153.9, 149.4, 125.5, 121.0, 106.0 ppm.

5 Example 6: 2-[4-(4-Isopropyl-phenoxy)-3,5-dimethyl-pyrazol-1-yl]-4-pyridin-2-yl-pyrimidine (Compound 97)

Step 1) Preparation of 3-(4-Isopropyl-phenoxy)-pentane-2,4-dione.

To a refluxing solution of 4-isopropylphenol (0.548 g, 4.02 mmol) and rhodium(II) acetate (ca. 15 mgs) in benzene (10 mL) was added a solution of 3-
10 diazo-pentane-2,4-dione (0.507 g, 4.02 mmol) in benzene (20 mL) over 40 minutes. The reaction mixture was allowed to cool to room temperature and was concentrated to afford a green oil. Flash chromatography over silica (ethyl acetate/hexanes) provided 0.371 g (39%) of the product as a white solid: ^1H NMR (CDCl_3) δ 14.4 (s, 1H), 7.15 (d, $J = 8.6$ Hz, 2H), 6.82 (d, $J = 8.6$ Hz, 2H), 2.87 (sept, $J = 7.0$ Hz,
15 1H), 2.04, (s, 6H), 1.23 (d, $J = 7.0$ Hz, 6H) ppm.

Step 2) Preparation of Compound 97.

To a solution of 3-(4-isopropyl-phenoxy)-pentane-2,4-dione (0.371 g, 1.58 mmol) in ethanol (20 mL) was added (4-pyridin-2-yl-pyrimidin-2-yl)-hydrazine (0.296 g, 1.58 mmol) and *p*-toluenesulfonic acid monohydrate (ca. 10 mgs). The solution was
20 heated to reflux and stirred for 14 hours. The reaction mixture was allowed to cool to room temperature, diluted with water (100 mL), and extracted with ethyl acetate (100 mL). The organic phase was washed with saturated sodium hydrogencarbonate solution (50 mL) and brine (50 mL), dried (magnesium sulfate), filtered, and concentrated to afford a yellow solid. Flash chromatography over silica
25 (methanol/methylene chloride) provided 0.421 g (69%) of the product as a light yellow solid: ^1H NMR (CDCl_3) δ 8.93 (d, $J = 5.1$ Hz, 1H), 8.75-8.74 (m, 1H), 8.46 (d, $J = 7.9$ Hz, 1H), 8.24 (d, $J = 5.1$ Hz, 1H), 7.88 (dt, $J = 7.9, 1.7$ Hz, 1H), 7.46-

7.42 (m, 1H), 7.14 (d, $J = 8.7$ Hz, 2H), 6.88 (d, $J = 8.7$ Hz, 2H), 2.88 (sept, $J = 6.9$ Hz, 1H), 2.69 (s, 3H), 2.23 (s, 3H), and 1.23 (d, $J = 6.9$ Hz, 6H) ppm. ^{13}C NMR (CDCl_3) δ 164.3, 160.2, 157.5, 156.4, 153.4, 149.7, 145.4, 142.5, 137.9, 137.2, 133.1, 127.4, 125.7, 122.0, 114.7, 114.0, 33.3, 24.1, 12.7, 11.3 ppm.

5

Example 7: 2-{4-[4-(4-Benzyl-piperazin-1-ylmethyl)-phenoxy]-3,5-dimethyl-pyrazol-1-yl}-4-pyridin-2-yl-pyrimidine (Compound 104)

Step 1) Preparation of {4-[3,5-Dimethyl-1-(4-pyridin-2-yl-pyrimidin-2-yl)-1H-pyrazol-4-yloxy]-phenyl}-methanol.

- 10 To a solution of 3-(4-acetoxymethyl-phenoxy)-pentane-2,4-dione (1.24 g, 4.69 mmol) in ethanol (30 mL) was added (4-pyridin-2-yl-pyrimidin-2-yl)-hydrazine (0.877 g, 4.69 mmol) and *p*-toluenesulfonic acid monohydrate (about 10 mgs). The reaction mixture was heated to reflux and stirred for 14 hours. The solution was allowed to cool to room temperature, and water (100 mL) was added. The resulting
- 15 precipitate was removed by filtration, washed with water, and dried to provide a white solid. This crude material was dissolved in 5:1 methanol/water (60 mL) and treated with excess potassium carbonate. The reaction mixture was stirred at room temperature for 2 hours and then diluted with water (100 mL). The precipitate that formed was isolated by filtration, washed with water, and dried to provide 1.53 g
- 20 (87%) of the product as a white solid. ^1H NMR analysis was consistent with the assigned structure.

Step 2) Preparation of 4-[3,5-Dimethyl-1-(4-pyridin-2-yl-pyrimidin-2-yl)-1H-pyrazol-4-yloxy]-benzaldehyde.

- To a solution of {4-[3,5-dimethyl-1-(4-pyridin-2-yl-pyrimidin-2-yl)-1H-pyrazol-4-yloxy]-phenyl}-methanol (1.53 g, 4.10 mmol) in chloroform (80 mL) was added the
- 25 Dess-Martin periodinane (2.36 g, 5.56 mmol). The reaction mixture was stirred at room temperature for 3 hours and then concentrated to provide a white paste. Flash

-42-

chromatography over silica (methanol/methylene chloride) provided 1.37 g (90%) the product as a white solid. ¹H NMR analysis was consistent with assigned structure.

Step 3) Preparation of Compound 104.

- 5 To a solution of 4-[3,5-dimethyl-1-(4-pyridin-2-yl-pyrimidin-2-yl)-1*H*-pyrazol-4-yloxy]-benzaldehyde (0.545 g, 1.47 mmol) and 1-benzylpiperazine (0.310 g, 1.76 mmol) in dichloroethane (10 mL) was added sodium triacetoxyborohydride (0.436 g, 2.06 mmol). The reaction mixture was stirred at room temperature for 17 hours. The solution was diluted with methylene chloride (50 mL) and washed with
- 10 saturated sodium hydrogencarbonate (2 x 30 mL) and brine (50 mL). The organic phase was dried (magnesium sulfate), filtered, and concentrated to afford a yellow oil. Flash chromatography over silica (2 M ammonia in methanol/methylene chloride) provided 0.183 g (23%) of the product as a white foam: ¹H NMR (CDCl₃)
- 15 δ 8.94-8.93 (m, 1H), 8.75 (d, *J* = 3.9 Hz, 1H), 8.46 (d, *J* = 7.9 Hz, 1H), 8.25 (d, *J* = 3.9 Hz, 1H), 7.90-7.87 (m, 1H), 7.46-7.43 (m, 1H), 7.31-7.21 (m, 7H), 6.89 (d, *J* = 8.5 Hz, 2H), 3.52 (s, 2H), 3.47 (s, 2H), 2.68 (s, 3H), 2.48 (br s, 8H), 2.21 (s, 3H) ppm.

Example 8: 2-[4-(4-Chloro-benzylsulfanyl)-3,5-dimethyl-pyrazol-1-yl]-4-pyridin-2-yl-pyrimidine (Compound 119)

- 20 Step 1) Preparation of 3-(4-Chloro-benzylsulfanyl)-pentane-2,4-dione.

- To a solution of 3-chloro-pentan-2,3-dione (1.94 g, 14.4 mmol) in ethylene glycol dimethyl ether (50 mL) was added sodium hydrogencarbonate (ca. 15 g) and 4-chlorobenzyl mercaptan (2.29 g, 14.4 mmol). The suspension was heated to reflux and stirred for 3 hours. The reaction mixture was allowed to cool to room
- 25 temperature, and water (200 mL) was added. The mixture was extracted with ethyl ether (100 mL x 2), and the combined organic extracts washed with brine (150 mL).

The organic phase was dried (magnesium sulfate), filtered, and concentrated to afford a yellow oil. Flash chromatography over silica (ethyl acetate/hexanes) provided 2.48 g (67%) of the product as a white solid: ^1H NMR (CDCl_3) δ 7.26 (d, $J = 8.4$ Hz, 2H), 7.06 (d, $J = 8.4$ Hz, 2H), 3.60 (s, 2H), 2.13 (s, 6H) ppm.

5 Step 2) Preparation of Compound 119.

To a solution of 3-(4-chloro-benzylsulfanyl)-pentane-2,4-dione (1.15 g, 4.48 mmol) in ethanol (60 mL) was added (4-pyridin-2-yl-pyrimidin-2-yl)-hydrazine (0.838 g, 4.48 mmol) and *p*-toluenesulfonic acid monohydrate (ca. 20 mgs). The solution was heated to reflux and stirred for 5 hours. The reaction mixture was
10 allowed to cool to room temperature, and water (200 mL) was added. The precipitate that formed was isolated by filtration and recrystallized from aqueous methanol to provide 1.36 g (74%) of the product as white needles: ^1H NMR (CDCl_3) δ 8.94-8.92 (m, 1H), 8.75-8.73 (m, 1H), 8.41 (d, $J = 7.9$ Hz, 1H), 8.28-8.26 (m, 1H), 7.91 (t, $J = 7.9$ Hz, 1H), 7.46-7.43 (m, 1H), 7.20 (d, $J = 8.0$ Hz, 2H),
15 6.89 (d, $J = 8.0$ Hz, 2H), 3.66 (s, 2H), 2.46 (s, 3H), 2.28 (s, 3H) ppm. ^{13}C NMR (CDCl_3) δ 164.6, 160.5, 157.4, 155.2, 153.5, 149.8, 147.3, 137.6, 137.2, 133.1, 130.6, 128.7, 126.1, 122.2, 114.7, 111.3, 40.1, 14.2, 12.6 ppm.

Example 9: 2-[4-(4-Chloro-phenylmethanesulfonyl)-3,5-dimethyl-pyrazol-1-yl]-4-pyridin-2-yl-pyrimidine (Compound 124)

20 To a suspension of 2-[4-(4-chloro-benzylsulfanyl)-3,5-dimethyl-pyrazol-1-yl]-4-pyridin-2-yl-pyrimidine (0.920 g, 2.26 mmol) and sodium hydrogencarbonate (4.10 g, 48.8 mmol) in acetone (125 mL) and water (45 mL) was added Oxone (3.47 g, 5.64 mmol). The reaction mixture was stirred at room temperature for 18 hours. The mixture was treated with excess sodium hydrosulfide, stirred for 15 minutes,
25 and extracted with ethyl acetate (3 x 50 mL). The combined organic phases were washed with brine (100 mL), dried (magnesium sulfate), filtered, and concentrated to afford a white solid. The crude product was triturated with hot methanol followed

by flash chromatography over silica (methanol/methylene chloride) to provide 0.230 g (23%) of the product as a white solid. ¹H NMR analysis was consistent with assigned structure.

Example 10: 6-[4-(4-Chloro-benzylsulfanyl)-3,5-dimethyl-pyrazol-1-yl]-
5 [2,2']bipyridinyl (Compound 126)

Step 1) Preparation of 4-(4-Chloro-benzylsulfanyl)-3,5-dimethyl-1*H*-pyrazole.

To a solution of crude 3-(4-chloro-benzylsulfanyl)-pentane-2,4-dione [prepared as described above from 3-chloro-pentan-2,3-dione (1.03 g, 7.63 mmol) and 4-chlorobenzyl mercaptan (1.21 g, 7.63 mmol)] in ethanol (60 mL) was added
10 hydrazine hydrate (0.713 mL, 22.9 mmol) and *p*-toluenesulfonic acid monohydrate (ca. 25 mg). The solution was heated to reflux and stirred for 15 hours. The reaction mixture was allowed to cool to room temperature, and water (300 mL) was added. The precipitate that formed was removed by filtration, washed with water, and dried to provide 1.38 g (72%) of the product as a white solid: ¹H NMR (CDCl₃)
15 δ 7.17 (d, *J* = 8.4 Hz, 2H), 6.92 (d, *J* = 8.4 Hz, 2H), 3.57 (s, 2H), and 2.02 (s, 6H) ppm. ¹³C NMR (CDCl₃) δ 137.2, 132.7, 130.3, 128.3, 105.2, 39.9, 10.7 ppm.

Step 2) Preparation of Compound 126.

To a solution of 4-(4-chloro-benzylsulfanyl)-3,5-dimethyl-1*H*-pyrazole (0.708 g, 2.80 mmol) in 2-methoxyethyl ether (7 mL) was added sodium hydride
20 (0.123 g (60% dispersion), 3.08 mmol). After stirring for 5 minutes, 6-bromo-[2,2']bipyridinyl (0.691 g, 2.94 mmol) was added, and the reaction mixture was heated to 100 °C for 16 hours. The reaction mixture was allowed to cool to room temperature, and water (100 mL) was added. The suspension was extracted with ethyl acetate (50 mL x 2), and the combined organic extracts washed with brine (100
25 mL). The organic phase was dried (magnesium sulfate), filtered, and concentrated to afford a tan solid. Flash chromatography over silica (methanol/methylene chloride)

followed by recrystallization from aqueous methanol provided 0.485 g (43%) of the product as a white solid: ^1H NMR (CDCl_3) δ 8.70-8.68 (m, 1H), 8.34 (d, $J = 7.9$ Hz, 1H), 8.29 (d, $J = 7.9$ Hz, 1H), 7.94 (t, $J = 7.9$ Hz, 1H), 7.86 (dt, $J = 2.0, 7.9$ Hz, 2H), 7.35-7.31 (m, 1H), 7.22 (d, $J = 8.3$ Hz, 2H), 6.99 (d, $J = 8.3$ Hz, 2H), 3.65 (s, 2H), 2.45 (s, 3H), 2.21 (s, 3H) ppm. ^{13}C NMR (CDCl_3) δ 155.7, 154.5, 153.8, 152.9, 149.5, 145.9, 139.7, 137.3, 137.2, 133.1, 130.7, 128.7, 124.2, 121.3, 118.5, 116.1, 110.0, 40.3, 13.6, 12.3 ppm.

Example 11: 4-[5-Benzyl-3-(4-chloro-phenyl)-[1,2,4]triazol-1-yl]-6-pyridin-2-yl-[1,3,5]triazin-2-ylamine (Compound 133)

10 Step 1) Preparation of *N*-(4-Chloro-thiobenzoyl)-2-phenyl-acetamide.

To a solution of 4-chlorothiobenzamide (0.519 g, 3.02 mmol) in acetone (5 mL) was added pyridine (0.367 mL, 4.53 mmol) and phenylacetyl chloride (0.480 mL, 3.63 mmol). The bright orange reaction mixture was heated to 55 °C for 1 hour. The reaction mixture was allowed to cool to room temperature, and water (20 mL) was added. The precipitate was removed by filtration, washed with water, and dried to provide 0.721 g (82%) of the product as a red solid: ^1H NMR (CDCl_3) δ 9.34 (br s, 1H), 7.50-7.22 (m, 9H), 3.96 (s, 2H) ppm.

Step 2) Preparation of Compound 133.

To a solution of *N*-(4-chloro-thiobenzoyl)-2-phenyl-acetamide (0.536 g, 1.85 mmol) and sodium acetate (ca. 0.25 g) in 1:1 acetic acid/1,4-dioxane (30 mL) was added 4-hydrazino-6-pyridin-2-yl-[1,3,5]triazin-2-ylamine (0.376 g, 1.85 mmol). The solution was heated to 90 °C and stirred for 15 hours. The reaction mixture was allowed to cool to room temperature, and water (100 mL) was added. The precipitate was removed by filtration, triturated with hot ethanol (300 mL), and dried to provide 0.489 g (60%) of the product as a white solid: ^1H NMR ($\text{DMSO}-d_6$) δ 8.78 (d, $J = 4.3$ Hz, 1H), 8.32 (d, $J = 6.1$ Hz, 2H), 8.22 (d, $J = 7.8$ Hz, 1H), 8.06 (d, J

= 8.5 Hz, 2H), 7.96 (dt, $J = 1.5, 7.8$ Hz, 1H), 7.62-7.57 (m, 3H), 7.36-7.26 (m, 2H), 7.24-7.22 (m, 2H), 7.19-7.16 (m, 1H), 4.90 (s, 2H) ppm.

Example 12: 2-[4-(4-Chloro-phenylsulfanyl)-3,5-dimethyl-pyrazol-1-yl]-4-pyridin-2-yl-[1,3,5]triazine (Compound 141)

5 Step 1) Preparation of 4-Pyridin-2-yl-[1,3,5]triazine-2-thiol.

To a stirred solution of picolamide (1.50 g, 12.3 mmol) in N,N-dimethylformamide (20 mL) was added *tert*-butoxybis(dimethylamino)methane (5.2 mL, 25 mmol). The mixture was refluxed for 2.5 hours and then concentrated. The resulting viscous brown oil was taken up in a 0.5 M solution of sodium ethoxide in
10 ethanol (50 mL, 25 mmol). Thiourea (0.72 g, 9.5 mmol) was added and the solution was refluxed for 2.5 hours. The reaction was concentrated and the residue partitioned between ethyl acetate and dilute aqueous sodium hydroxide solution. The aqueous layer was washed twice with ethyl acetate and neutralized with 1 N aqueous hydrochloric acid. The resulting precipitate was filtered off and rinsed with
15 water. Vacuum oven drying afforded 0.72 g (40%) of crude product as a fine brown solid. This material was used without further purification in the next step.

Step 2) Preparation of 2-Methylsulfanyl-4-pyridin-2-yl-[1,3,5]triazine.

To a stirred suspension of the product of step 1 (0.707 g, 3.72 mmol) in acetone (36 mL) was added sodium carbonate (0.79 g, 7.5 mmol) followed by
20 iodomethane (0.35 mL, 5.6 mmol). After overnight stirring, the reaction was filtered free of undissolved solid and concentrated to afford product as light brown solid: ^1H NMR (DMSO- d_6) δ 9.13 (s, 1H), 8.84-8.76 (m, 1H), 8.55-8.43 (m, 1H), 8.11-8.00 (m, 1H), 7.69-7.60 (m, 1H), 2.63 (s, 3H) ppm.

Step 3) Preparation of (4-Pyridin-2-yl-[1,3,5]triazin-2-yl)-hydrazine.

-47-

A stirred solution of the product of step 2 (0.763 g, 3.74 mmol) and hydrazine hydrate (0.22 mL, 3.9 mmol) in ethanol (8 mL) was heated at reflux for 30 minutes. The reaction was cooled in an ice bath and the precipitate was filtered off. Vacuum oven drying afforded crude product as light brown solid. This material was
5 used without further purification in the next step.

Step 4) Preparation of Compound 141.

A stirred suspension of the product of step 3 (0.333 g, 1.77 mmol) and 3-(4-chloro-phenylsulfanyl)-pentane-2,4-dione (0.473 g, 1.95 mmol) in *n*-propanol was heated at 95°C overnight. The reaction solution was concentrated directly onto silica
10 and flash chromatographed (methylene chloride/methanol) to afford a dirty yellow solid. This material was further purified by trituration with diethyl ether to afford 0.056 g (8%) of product as a white solid: ¹H NMR (DMSO-d₆) δ 9.44 (s, 1H), 8.94-8.77 (m, 1H), 8.68-8.43 (m, 1H), 8.18-7.96 (m, 1H), 7.77-7.57 (m, 1H), 7.45-7.24 (m, 2H), 7.20-6.98 (m, 2H), 2.83 (s, 3H), 2.20 (s, 3H) ppm.

15 Example 13: 4-(4-Benzyl-3,5-dimethyl-pyrazol-1-yl)-6-pyridin-2-yl-[1,3,5]triazin-2-ylamine (Compound 142)

Step 1) Preparation of 3-Benzylpentane-2,4-dione.

To a solution of sodium ethoxide (freshly prepared from sodium hydride (0.474 g, 19.8 mmol) and anhydrous ethanol (50 mL)) was added pentane-2,4-dione
20 (6.00g, 60.0 mmol). The resulting mixture was heated to 50 °C while a solution of benzyl bromide (3.42 g, 20 mmol) in ethanol (20 mL) was added over 30 minutes. The reaction was then heated to reflux. After 2 hours, the mixture was concentrated, and the residue was dissolved in ethyl acetate, washed three times with water, once with brine, dried (magnesium sulfate) and concentrated to afford an oil. Flash
25 chromatography over silica (ethyl acetate/dichloromethane) afforded 3.13 g (82%) of a colorless oil. The proton spectrum showed the product exists as equal parts of the

keto and enol tautomers: ^1H NMR (CDCl_3) δ 16.81 (s, 0.5H), 7.13-7.32 (m, 5H), 4.00 (t, $J = 7.6$ Hz, 0.5H), 3.65 (s, 1H), 3.14 (d, $J = 7.6$ Hz, 0.5H), 2.06-2.13 (m, 6H) ppm.

Step 2) Preparation of Compound 142.

5 A suspension of 4-hydrazino-6-pyridin-2-yl-[1,3,5]triazin-2-ylamine (5.25 g, 25.6 mmol), 3-benzylpentane-2,4-dione (5.0 g, 28.4 mmol), and *p*-toluenesulfonic acid monohydrate (0.475 g, 2.50 mmol) in dimethylsulfoxide (50 mL) was heated to 75 °C for 20 hours. The reaction mixture was allowed to cool to room temperature, and the suspension was dissolved in methylene chloride (700 mL). The solution was
10 washed three times with water, once with brine, dried (magnesium sulfate), and concentrated to afford a moist solid. The crude product was suspended in ethyl acetate (50 mL), heated to reflux, allowed to cool to room temperature, and filtered (repeated three times). The final product was obtained as a white solid containing 3 % dimethylsulfoxide by weight: 8.63 g (90%). ^1H NMR (CDCl_3) δ 8.84-8.53 (m,
15 1H), 8.48 (d, $J = 7.9$ Hz, 1H), 7.86-7.91 (m, 1H), 7.46-7.50 (m, 1H), 7.13-7.30 (m, 5H), 6.61 (br s, 1H), 6.40 (br s, 1H), 3.83 (s, 2H), 2.79 (s, 3H), 2.24 (s, 3H) ppm. MS (ESI) m/z 358 ($\text{M} + \text{H}^+$).

Example 14: 4-[3,5-Dimethyl-4-(3-phenyl-propyl)-pyrazol-1-yl]-6-pyridin-2-yl-[1,3,5]triazin-2-ylamine (Compound 144)

20 Step 1) Preparation of 3-(3-Phenyl-propyl)-pentane-2,4-dione.

A mixture of pentane-2,4-dione (3.87 g, 38.6 mmol), 3-phenyl-1-iodopropane (3.18g, 12.9 mmol) and anhydrous potassium carbonate (1.7 g, 12.3 mmol) in acetone (7.5 mL) was heated to reflux for 24 hours. The room temperature reaction mixture was filtered, and the filter cake washed with acetone
25 (25 mL x 3). The combined filtrates were concentrated, and the residue was partitioned between ethyl acetate and water. The organic layer was washed with

water and brine, dried (magnesium sulfate), and concentrated to an oil. Flash chromatography over silica (ethyl acetate/hexanes) afforded 1.13 g (42%) of a colorless oil. The proton spectrum in deuteriochloroform showed the product exists as a mixture of keto and enol tautomers.

5 Step 2) Preparation of Compound 144.

A solution of 4-hydrazino-6-pyridin-2-yl-[1,3,5]triazin-2-ylamine (6.17 g, 30.4 mmol), 3-(3-phenyl-propyl)-pentane-2,4-dione (6.66 g, 30.4 mmol), and *p*-toluenesulfonic acid monohydrate (0.05g, 0.26 mmol) in dimethylsulfoxide (200 mL) was heated 75 °C for 18 hours. The solution was allowed to cool to room temperature then diluted with ethyl acetate (700 mL). The resulting solution was washed 3 times with water, once with brine, dried (magnesium sulfate) and concentrated to afford a cream colored solid. The solid was triturated in boiling ethyl acetate (200 mL), and the suspension allowed to cool to room temperature. After 2 hours, the product was collected by filtration and dried *in vacuo* to yield 5.87 g (50%) of a white solid: ¹H NMR (CDCl₃) δ 8.84-8.54 (m, 1H), 8.49 (d, *J*=7.9 Hz, 1H), 7.87-7.91 (m, 1H), 7.46-7.50 (m, 1H), 7.18-7.32 (m, 5H), 6.44 (br s, 1H), 6.25 (br s, 1H), 2.73 (s, 3H), 2.65-2.73 (m, 2H), 2.44-2.49 (m, 2H), 2.31 (s, 3H), 1.79-1.86 (m, 2H) ppm. MS (ESI) *m/z* 386(M+H⁺).

Example 15: 4-(4-Chloro-phenylsulfanyl)-5-methyl-2-(4-pyridin-2-yl-pyrimidin-2-yl)-2,4-dihydro-pyrazol-3-one (Compound 147)

Step 1) Preparation of 2-(4-Chloro-phenylsulfanyl)-3-oxo-butyric acid ethyl ester.

To a solution of ethyl 2-chloroacetoacetate (1.58 g, 9.61 mmol) in ethylene glycol dimethyl ether (30 mL) was added sodium hydrogencarbonate (ca. 15 g) and 4-chlorothiophenol (1.39 g, 9.61 mmol). The suspension was heated to reflux and stirred for 3 hours. The reaction mixture was allowed to cool to room temperature, and water (100 mL) was added. The mixture was extracted with ethyl ether (75

mL), and the organic extract was washed with brine (150 mL). The organic phase was dried (magnesium sulfate), filtered, and concentrated to afford a yellow oil. Flash chromatography over silica (ethyl acetate/hexanes) provided 1.44 g (55%) of the product as a colorless oil: ^1H NMR (CDCl_3) δ 13.8 (s, 1H), 7.26-7.21 (m, 2H), 7.06-7.04 (m, 2H), 4.21 (q, $J = 7.1$ Hz, 2H), 2.33 (s, 3H), 1.20 (t, $J = 7.1$ Hz, 3H).

Step 2) Preparation of Compound 147.

To a solution of 2-(4-chloro-phenylsulfanyl)-3-oxo-butyric acid ethyl ester (4.13 g, 15.1 mmol) in n-propanol (85 mL) was added (4-pyridin-2-yl-pyrimidin-2-yl)-hydrazine (2.84 g, 15.1 mmol) and *p*-toluenesulfonic acid monohydrate (ca. 25 mgs). The solution was heated to reflux and stirred for 15 hours. The reaction mixture was allowed to cool to room temperature, diluted with water (400 mL), and extracted with ethyl acetate (250 mL). The organic phase was washed with water (200 mL) and brine (200 mL), dried (magnesium sulfate), filtered, and concentrated to afford a brown oil. The crude material was triturated with diethyl ether, and the solid that formed was isolated by filtration. Further trituration with a minimum amount of hot methanol provided the product as a tan solid. ^1H NMR (CDCl_3) δ 12.8 (br s, 1H), 8.85 (d, $J = 5.3$ Hz, 1H), 8.77 (d, $J = 4.3$ Hz, 1H), 8.44 (d, $J = 7.8$ Hz, 1H), 8.29 (d, $J = 5.3$ Hz, 1H), 7.92 (dt, $J = 1.5, 7.8$ Hz, 1H), 7.51 - 7.48 (m, 1H), 7.19 (d, $J = 8.6$ Hz, 2H), 7.09 (d, $J = 8.6$ Hz, 2H), 2.32 (s, 3H). ^{13}C NMR (CDCl_3) δ 165.1, 159.1, 158.8, 157.1, 156.8, 152.0, 137.7, 136.6, 131.3, 129.2, 127.3, 126.8, 122.7, 114.6, 88.9, 13.3.


Example 16: 2-[4-(4-Chloro-phenylsulfanyl)-5-methoxy-3-methyl-pyrazol-1-yl]-4-pyridin-2-yl-pyrimidine (Compound 148)

To a solution of 4-(4-chloro-phenylsulfanyl)-5-methyl-2-(4-pyridin-2-yl-pyrimidin-2-yl)-2, 4-dihydro-pyrazol-3-one (0.344g, 0.869mmol) in acetone (10mL) was added potassium carbonate (ca. 3g) and iodomethane (0.130g, 0.912 mmol). The suspension was heated to reflux and stirred for 2 hours. The suspension was

































-51-





























allowed to cool to room temperature, and water (100mL) was added. The mixture was extracted with diethyl ether (50mL), and the organic phase was washed with brine (50 mL), dried (magnesium sulfate), filtered and concentrated to afford a white solid. Flash chromatography over silica (methylene chloride/2M ammonia in
5 methanol, two columns) provided 0.007g (2%) of the product as a white solid. ¹H NMR analysis was consistent with the assigned structure.

































Table I:



































































Cpd No.	Method Used	R ₁₁	R ₁₀	R ₉	R ₈	R ₁₂
1	1					H
2	1					H
3	1					H
4	1					H
5	1					H
6	1					H
7	1					H

































8	1					H
9	1					H
10	1					H
11	1					H
12	1					H
13	1					H
14	1					H
15	1					H
































16	1					H
17	1					H
18	1					H
19	1					H
20	1					H
21	1					H
22	1					H

































23	1					H
24	1					H
25	1					H
26	4					CH ₃
27	1					H
28	1					H
29	1					H
30	1					H

































31	1					H
32	1					H
33	1					H
34	1					H
35	1					H
36	1					H
37	1					H
38	1					H

































39	1					H
40	1					H
41	1					H
42	1					H
43	1					H
44	1					H
45 ^a	3					H
46 ^a	3					H

























47 ^b	3					H
48 ^b	3					H
49 ^c	1					H
50 ^c	1					H
51	2					H
52	2					H
53	2					H
54	2					H

55	2					H
56	2					H
57	2					H
58	2					H
59	2			H,H		H
60	2					H
61	2					H
62	2					H

63	2					H
64	2					H
65	2					H
66	2					H
67	2					H
68	2					H
69	2					H
70	2					H

71	2					H
72	2					H
73	2					H
74	2					H
75	2					H
76	2					H
77	2					H
78	2					H

79	2					H
80	2					H
81	2					H
82	2					H
83	2					H
84	2					H
85	2					H
86	2					H

87	2					H
88	2					H
89	2					H
90	2					H
91	2					H
92	2					H


^aCompound 45 was prepared at the (*R*)-enantiomer. Compound 46 was prepared as the (*S*)-enantiomer. Enantiomeric excesses were determined to be 91.4% and 91.8% respectively.

^bCompound 47 was prepared at the (*R*)-enantiomer. Compound 48 was prepared as the (*S*)-enantiomer. Enantiomeric excesses were determined to be 100% (within detection limits) and >95% respectively.

^cCompounds 49 and 50 are the (-) and (+) enantiomers, respectively, of compound 44.

Preparative chiral HPLC was used for the resolution (ChiralPack OD, 2 x 25 cm; eluant: 40/60 carbon dioxide/acetonitrile; detection 260 nm).

Table II




Cpd No.	Ex. No.	R ₁₉	R ₄	R ₃	n	X	W	V
5	93	H	H	CH ₃	0	O	N	CH
	94	4-Cl	H	CH ₃	0	O	N	CH
	95	4-F	H	CH ₃	0	O	N	CH
	96	4-CH ₃ O	H	CH ₃	0	O	N	CH
	97	4-(CH ₃) ₂ CH	H	CH ₃	0	O	N	CH
10	98	4-CF ₃	H	CH ₃	0	O	N	CH
	99	4-CF ₃ O	H	CH ₃	0	O	N	CH
	100	4-(CH ₂ OH)	H	CH ₃	0	O	N	CH
	101	4-CHO	H	CH ₃	0	O	N	CH
	102	4-(N-morpholino)CH ₂	H	CH ₃	0	O	N	CH
15	103	4-(N-tetrahydroisoquinolino)CH ₂	H	CH ₃	0	O	N	CH
	104	4-(4-Benzyl-piperazin-1-yl)CH ₂	H	CH ₃	0	O	N	CH
	105	4-(4-(2-fluoro-phenyl)piperazin-1-yl)CH ₂	H	CH ₃	0	O	N	CH
	106	4-Cl	H	CH ₃	0	O	N	N
	107	4-Cl	H	CH ₃	1	O	N	CH
20	108	H	H	CH ₃	0	S	N	CH

5	109	8	2-Cl	H	CH ₃	0	S	N	CH
	110	8	3-Cl	H	CH ₃	0	S	N	CH
	111	8	4-Cl	H	CH ₃	0	S	N	CH
	112	8	4-F	H	CH ₃	0	S	N	CH
	113	8	4-CH ₃ O	H	CH ₃	0	S	N	CH
10	114	8	3,4-diCl	H	CH ₃	0	S	N	CH
	115	8	4-Cl	4-CH ₃	CH ₃	0	S	N	CH
	116	8	4-Cl	5-CH ₃	CH ₃	0	S	N	CH
	117	8	H	H	CH ₃	1	S	N	CH
	118	8	2-Cl	H	CH ₃	1	S	N	CH
15	119	8	4-Cl	H	CH ₃	1	S	N	CH
	120	8	4-F	H	CH ₃	1	S	N	CH
	121	8	4-CH ₃	H	CH ₃	1	S	N	CH
	122	8	4-CH ₃ O	H	CH ₃	1	S	N	CH
	123	8	4-CF ₃ O	H	CH ₃	1	S	N	CH
20	124	9	4-Cl	H	CH ₃	1	SO ₂	N	CH
	125	10	4-Cl	H	CH ₃	0	O	C	CH
	126	10	4-Cl	H	CH ₃	1	S	C	CH
	147	15	4-Cl	H	OH	0	S	N	CH
	148	16	4-Cl	H	OCH ₃	0	S	N	CH

Table III


-66-



	Cmp No.	Method Used	R ₂₁	R ₂₂
5	127	11	2,4-diClPh	4-ClPhCH ₂
	128	11	2,4-diFPh	4-ClPhCH ₂
	129	11	4-t-BuPh	4-ClPhCH ₂
	130	11	4-ClPh	4-ClPhCH ₂
	131	11	2,4-diFPh	4-(CH ₂ O)PhCH ₂
10	132	11	4-ClPh	3-(CH ₂ O)PhCH ₂
	133	11	4-ClPh	PhCH ₂
	134	11	4-ClPh	c-C ₆ H ₁₁
	135	11	4-ClPh	c-(C ₆ H ₉)CH ₂ CH ₂
	136	11	4-ClPh	c-C ₆ H ₆
15	137	11	4-ClPh	(CH ₃) ₂ CCH ₂
	138	11	4-ClPh	3,4-di(CH ₂ O)PhCH ₂

Table IV

-67-



	Cpd No.	Method Used	R ₂₃	R ₂₄	n	X
5	139	8	4-Cl	NH ₂	0	S
	140	9	4-Cl	NH ₂	0	SO ₂
	141	12	4-Cl	H	0	S
	142	13	H	NH ₂	0	C
	143	13	4-Cl	NH ₂	1	C
10	144	14	H	NH ₂	2	C
	145	13	4-Cl	NH ₂	1	C
	146	14	H	NH ₂	2	C

II BIOLOGICAL RESULTS

Example 17: In Vitro Activity of Compounds in TNF- α and VCAM Assays

Description of Table V: The results shown in Table V were obtained using the TNF- α assays performed as described in the protocols entitled "Method for TNF High Throughput Screen" and "Method for VCAM High Throughput Screen", which are detailed below. These assays measure a cellular response to TNF- α stimulation, and the data published in the table are expressed as percentage inhibitions. Percentage inhibition is computed on a scale of 0 to 100% where 0 percent inhibition means the compound has no effect in the assay (the response is indistinguishable from TNF- α treatment alone), and 100% inhibition means that when the compound is present at the stated concentration, the response in the assay is the same as if TNF- α were not added. These two assays measure two different outcomes of TNF- α treatment, cell death and Vascular Cell Adhesion Molecule (VCAM) expression, and capture two of TNF- α 's many biological effects. TNF- α induced apoptosis is a mechanism by which TNF- α production during immune/inflammatory responses can lead to direct cellular and tissue damage. TNF- α VCAM expression in endothelial cells is a response that allows leukocytes to leave the blood and enter a site of immune/inflammatory response. Both of these activities could lead to pathology in immune/inflammatory diseases. In the first column of the table, the percentage inhibition of the TNF- α driven apoptosis assay when the compounds are tested at 1 μ M concentration is presented. The second column in the table lists the IC₅₀ for the compound, a standard measure using a dilution series in the assay, and fitting a sigmoidal curve to the resulting dose-response curve. The IC₅₀ is the concentration (derived analytically from the sigmoidal function) at which the compound would inhibit 50% of the response in the assay. The third and fourth column list the corresponding analysis for the TNF- α driven VCAM assay. In Table VI, data corresponding to the data in column 1 of Table V is shown, with human cells replacing the mouse cells used in the high throughput screen. It will be understood that the table demonstrates that a class of compounds can have its best activity in one assay or the other, or that there can be crossover such that the compounds can have activity to some degree in both assays. While not being bound by theory, we

believe this data shows that the compounds can have a modulating effect in more than one activity. As a consequence, the compounds will have use in treating more than one TNF- α mediated condition, and may be selected according to the manner in which the condition manifests. This corresponds to the use of these compounds in

5 the treatment of human disease.

TNF High Throughput Screen.

Protocol:

1. Plate 4×10^4 L929 cells in 100 μ l of complete EMEM in Costar 96 well plates in PM.
- 10 2. The next day, pretreat with compound, inhibitor and control vehicle for 2 hours.
3. After 2 hours pretreatment, add 10 μ l 5ng/ml human TNF- α and actinomycin D (40 μ g/ml final concentration).
4. Incubate overnight.
- 15 5. In AM, remove supernatant.
6. Wash on a plate washer (0.9% NaCl).
7. Add 100 μ l Crystal Violet 0.1% in 20% ETOH.
8. Incubate at RT for 10 minutes.
9. Wash on plate washer.
- 20 10. Air dry wells at 37°C.
11. Add 100 μ l of methanol to each well.
12. Shake plates on orbital shaker and read at 595 nm on plate reader.

VCAM High Throughput Screen:

Protocol:

- 25 1. Plate Primary human umbilical vein endothelial cells at 1.8×10^4 cells/well in a 96-well tissue culture plate.
2. Return Plates to 37°C incubator for 48-72 hours before assay.

-70-

3. On assay day, aspirate wells and add 180 μ L of endothelial growth cell medium (EGM) to each well.
4. Add Compound to each well.
5. Shake plates for 3 minutes.
- 5 6. Incubate plate for 1 hour at 37°C.
7. Add Tumor Necrosis Factor (TNF) at 1.0 ng/ml final concentration.
8. Shake plates for 3 minutes.
9. Incubate plate for 2 hours at 37°C.
10. Remove plate from incubator and wash plate 3 times with phosphate buffered saline (PBS) using a plate washer.
- 10 11. Add anti-VCAM antibody at 0.5 μ g/ml final concentration.
12. Incubate at 4°C overnight.
13. Wash 3 times with PBS on plate washer.
14. Add goat anti-mouse horseradish peroxidase conjugate.
- 15 15. Incubate at room temperature for one hour.
16. Wash 3 times with PBS on plate washer.
17. Add TMB to each well for 15 minutes at room temperature.
18. Stop reaction with 100 μ L of 2 N sulfuric acid.
19. Read absorbance at 450 nm.

20

Table V: In Vitro Activity of Compounds in TNF- α and VCAM Assays

Cpd. No.	TNF α %Inh. @ 1 μ M	TNF α IC ₅₀ (μ M)	VCAM %Inh. @12.5 μ M	VCAM IC ₅₀ (μ M)
1		2.64		
25 2		5.65		
3		1.29		

-71-

	4		0.941		
	5		0.872		
	6		1.06		
	7		4.01		
5	8		1.62		
	9		0.496		
	10		3.68		
	11		2.89		
10	12		5.15		
	13		1.37		
	14		1.79		
	15		0.386		
	16		1.32		
	17		1.5		
15	18		2.68		
	19		0.457		
	20		0.299		
	21		0.913		
	22		3.85		
20	23		2.58		
	24		1.03		
	25		1.58		
	26		3.36		
	27		2.72		
25	28		5.27		
	29		3.72		
	30		2.36		
	31		3.35		
	32		2.33		
30	33		2.35		
	34		1.54		
	35		2.64		
	36		2.15		
	37		2.36		
35	38		2.79		
	39		1.32		
	40		1.74		
	41		6.49		
	42		0.779		16
40	43		2.98		
	44		0.41		24.4
	45		3.34		

-72-

	46		4.37		
	47		5.58		
	48		4.05		
	49		1.53		
5	50		0.39		
	51		3.56		
	52		3.49		
	53		7.13		
	54		6.03		
10	55		7.64		
	56		6.63		
	57	56			
	58	50			
	59	51			
15	60	50			
	61	51			
	62	59			
	63	55			
	64	63			
20	65	52			
	66	51			
	67	52			
	68	57			
	69	61			
25	70	58			
	71	57			
	72	57			
	73	76			
	74	55			
30	75	67			
	76	56			
	77	54			
	78	60			
	79	62			
35	80	53			
	81	50			
	82	55			
	83	56			
	84	52			
40	85	52			
	86	53			
	87	50			

-73-

	88	53		
	89	56		
	90	51		
	91	71		
5	92	52		
	93		1.21	
	94		0.029	
	95		0.2	
	96		0.046	
10	97		0.397	
	98		0.058	
	99		0.177	
	100		3.46	
	101		17.9	
15	102		0.431	
	103		0.709	
	104		0.05	
	105		0.089	
	106		0.96	
20	107		47	
	108		2.28	
	109		0.63	
	110		6.34	
	111		0.132	
25	112		0.309	
	113		0.498	
	114		1.44	
	115		19.4	
	116		355	
30	117			16
	118		72	
	119			7.95
	120			13.2
	121			9.93
35	122			11.9
	123			7.1
	124			13.4
	125		0.023	
	126		37.9	2.52
40	127		2.2	
	128		20.6	8.67
	129		19.5	

-74-

	130		8.96		
	131		12.8		
	132		1.39		
	133		19.5		
5	134		0.526		
	135		0.275		
	136		1.44		
	137		9.67		
	138	54 ^a			
10	139	62 ^b			
	140		1.77		
	141		0.0011		
	142		0.152		
	143		6.06		
15	144		0.015		
	145		2.5		
	146		0.197		
	147		0.734		
	148		32.1		

20

^a Inhibition measured at 2 μ M^b Inhibition measured at 25 μ M

Table VI. Compound 44 can inhibit the TNF- α induced apoptosis in primary human fibroblasts

		OD 595nm	Standard Deviation	% Rescue
	DMSO	0.189	0.005	N/A
25	TNF + DMSO	0.091	0.006	N/A
	2 μ M Compound 44	0.109	0.008	18.4
	4 μ M Compound 44	0.147	0.033	57.1
	8 μ M Compound 44	0.198	0.002	109.2

Normal human dermal fibroblasts were seeded into 96-well plates at 3 x 10⁴ cells/well. Cells were pretreated for 2 hours with 2 μ g/ml Actinomycin D and either

30

Compound 44 or DMSO as a vehicle control. Cells were then exposed to 2ng/ml TNF- α for 24 hours and stained with crystal violet/ethanol. Crystal violet was solubilized with methanol and absorbance at 595 nm was read on a microtiter plate reader.

5 Example 18: In Vivo Activity of Compounds in Sepsis and IBD Models and Pharmacokinetic Parameters

Sepsis Model:

Sepsis was induced in the C57/BL mouse by the intravenous injection of 20
10 ng of lipopolysaccharide/animal plus 20 mgs d-galactosamine/animal. Inhibition was measured as the prevention of mortality over a three day period. Compounds were administered intraperitoneally in 10% cremophore/10% ethanol/80% normal saline one hour before induction.

Inflammatory Bowel Disease Model:

15 Groups of three male rats weighing 150 +/-10g and fasted for 24 hours were used. Distal colitis was induced by intracolonic instillation of 0.5ml/rat DNBS (2,4-dinitrobenzene sulfonic acid, 60 mg/ml in ethanol 30%) after which air (2 ml) was gently injected through the cannula to ensure that the solution remains in the colon. A test compound was administered orally 24 and 2 hours before DNBS-instillation
20 and then daily for 5 days in a total of 7 doses. The animals were sacrificed 24 hours after the final dose of test compound administration and each colon was removed and weighed.

Table VII: In Vivo Activity of Compounds in Sepsis and IBD Models and Pharmacokinetic Parameters

	<u>Sepsis:</u> Murine LPS-d-galactosamine model #	<u>Inflammatory</u> <u>Bowel Disease:</u> Rat DNBS model*	<u>Pharmacokinetic</u> <u>Parameters</u>	
	% inhibition	% inhibition of colonic weight gain	$t^{1/2}$	% oral bioavailability
44	33+/- 1 @ 200 mg/kg (2 exp.)	44 +/- 8 @ 100 mg/kg (2 exp.)	2.3	26
119	33 @ 200 mg/kg	51% @ 100 mg/kg	2.3	Undetectable
94	Not active below 200mg/kg	18% @ 20 mg/kg	10.9	Undetectable
43	33 @ 10 mg/kg	31% @ 50 mg/kg	3.6	<1

5 Example 19: In Vivo Activity of Compounds in Experimental Allergic
Encephalitis (Murine Model of Multiple Sclerosis)

SJL mouse strain were immunized subcutaneously with proteolipid peptide
amino acid residues 139-151 in Complete Freund's Adjuvant. Pertussis toxin was
injected intravenously on day 1 and 3 post-induction. Weight loss occurred at day 7
10 post-immunization with paralysis ensuing between days 9 and 14 post-
immunization.

Table VIII: In Vivo Activity of Compounds in Experimental Allergic
Encephalitis (Murine Model of Multiple Sclerosis)

15

Compound	Dosing regimen ¹	Ratio of activity disease severity score (ADSS) ⁴ at last day of dosing for treated vs. control	% inhibition of disease ⁵	% Survival ⁷ of treated vs. %survival of controls
----------	-----------------------------	--	---	---

-77-

5	44	15	1.9/3.35	44.0%	75 vs. 50
	Induction regimen	mgs/kg/day/i.p. ² from day 1-21		P ⁶ =0.06	
		75 mgs/kg/day/i.p. from day 1-21	0.8/3.35	75.7%	100 vs. 50
				p=0.001	
	44	15 mgs/kg/day/i.p. from day 7-21	2.6/4.6	43.5%	70 vs. 11
	Therapeutic regimen			p=0.08	
10		75 mgs/kg/day/i.p. from day 7-21	1.1/4.6	76.1%	100 vs. 11
				p=0.001	
	44	75 mgs/kg/day/i.p. from day 7-28	2.2/4.4	50%	89 vs 20
	Oral therapeutic regimen			p=0.002	
	94	75 mgs/kg/day/i.p. from day 1-18	2.15/3.53	39.1%	80 vs 67
	Induction Regimen			p ⁶ =0.045	

- Compounds solublized in 10 % cremophore and 10% ethanol 80% normal saline.
- Compounds administered intraperitoneally.
- Compound administered orally.
- ADSS is activity disease severity score where >1 is significant disease. Paralysis starts at the tail and progresses towards head of the mouse where a limp tail is a 1 and a complete paralysis is a 5. Animals are euthanized at greater than or equal to a score of 4.
- % inhibition of ADSS = (average test ADSS - average control ADSS/control) x 100.
- P values calculated using Student's T Test.
- Survival indicates those animals remaining post-euthanasia.


20 EQUIVALENTS

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

CLAIMS

What is claimed is:

1. A compound of Formula I,



5

and physiological salts thereof, wherein:

n is 0, 1 or 2;

X is O, CH₂, S or SO₂;

R₁ is H or NH₂;


10 R₂ and R₃ are each, independently, -H, -OH, a substituted or unsubstituted alkyl, or a substituted or unsubstituted alkoxy;

R₄ is, -H or a substituted or unsubstituted alkyl;

V, W and Z are each, independently, N or CH; and


Y is selected from the group consisting of substituted and unsubstituted phenyl, substituted and unsubstituted heterocyclyl.

2. The compound of Claim 1, wherein Y is a phenyl group which has one or more substituents independently selected from the group consisting of halogen, linear or branched C₁-C₄-alkoxy, trifluoromethoxy, dioxymethylene, hydroxyalkyl, trifluoromethyl, HC(O)-, linear or branched C₁-C₄-alkyl, heterocyclyl and substituted or unsubstituted heterocycloalkylalkyl.
- 5
3. The compound of Claim 2, wherein Y is a phenyl group which has one or more substituents selected from the group consisting of fluoro, chloro, methoxy, morpholyl, N-morpholinomethyl, tetrahydroisoquinolyl, tetrahydroisoquinolinomethyl, 4-(4-benzyl-piperazin-1-yl)methyl, 4-(4-(2-fluoro-phenyl)piperazin-1-yl)methyl, and isopropyl.
- 10
4. The compound of Claim 1, wherein Y is selected from the group consisting of pyridyl, furyl, and pyrrolidyl.
5. A compound represented by the following structural formula:




- 15 6. A compound represented by the following structural formula:

-80-



7. A compound of Formula II,



-81-

and physiological salts thereof, wherein,

R₅ is substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted cycloalkylalkyl;

R₆ is -H or -NR₁₃R₁₄;

5 R₇ is substituted or unsubstituted phenyl; and


R₁₃ and R₁₄ are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl; or

10 R₁₃ and R₁₄ together with the nitrogen to which they are attached are a heterocycloalkyl.

8. The compound of Claim 7, wherein R₅ is substituted or unsubstituted benzyl.
9. The compound of Claim 8, wherein R₅ is benzyl having one or more substituents independently selected from the group consisting of halogen, linear C₁-C₄-alkoxy and branched C₁-C₄-alkoxy.
- 15 10. The compound of Claim 9, wherein R₅ is benzyl having one or more substituents independently selected from the group consisting of chloro and methoxy.
11. The compound of Claim 7, wherein R₅ is C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-C₁-C₄-alkyl or substituted or unsubstituted phenyl-C₂-C₄-alkyl.
- 20 12. The compound of Claim 11, wherein R₅ is selected from the group consisting of 2-phenethyl, cyclohexyl and cyclopentylethyl.
13. The compound of Claim 7, wherein R₇ is phenyl having one or more substituents independently selected from the group consisting of halogen, linear C₁-C₆-alkyl, branched C₁-C₆-alkyl, cyclic C₃-C₆-alkyl and
25 trifluoromethyl.

14. The compound of Claim 13, wherein R_7 is phenyl having one or more substituents independently selected from the group consisting of fluoro, chloro, linear C_1 - C_4 -alkyl, and branched C_1 - C_4 -alkyl.
15. A compound of Formula III,

5



and physiological salts thereof, wherein:

R_8 and R_{12} are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, substituted or unsubstituted aralkyl or a substituted or unsubstituted heteroaralkyl;

10

R_9 is -H, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl, a substituted or unsubstituted heteroaryl or a substituted or unsubstituted heteroaralkyl;

15

R_{10} is substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaralkyl, or a substituted or unsubstituted heterocycloalkylalkyl; and

20

R_{11} is a substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl, a substituted or unsubstituted cycloalkylalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted heteroaralkyl, a substituted or unsubstituted benzophenone, or a substituted or unsubstituted cycloalkylalkyl.

16. The compound of Claim 15, wherein one of R_8 or R_{12} is -H and the other is substituted or unsubstituted phenyl, phenyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -

alkyl, linear C₁-C₁₂-alkyl, branched C₁-C₁₂-alkyl, cyclic C₃-C₁₂-alkyl, or dicycloalkyl-C₁-C₄-alkyl.


17. The compound of Claim 16, wherein one of R₈ or R₁₂ is -H and the other is phenyl, phenyl-C₁-C₄-alkyl, or diphenyl-C₁-C₄-alkyl wherein the phenyl group or phenyl groups bear one or more substituents independently selected from the group consisting of C₁-C₄-alkoxy, C₁-C₄-alkyl and cyano.
18. The compound of Claim 17, wherein the phenyl group or phenyl groups bear one or more substituents independently selected from the group consisting of methoxy, methyl, ethyl and cyano.
19. The compound of Claim 15, wherein R₈ is selected from the group consisting of 2,2-diphenylethyl, 2-(4-ethylphenyl)ethyl, benzyl, diphenylmethyl, 1,2-diphenylethyl, 3,3-diphenylpropyl, 3,4,5-trimethoxybenzyl, 2,4,4-trimethylisopentyl, 2-(4-methoxyphenyl)ethyl, 2-cyclopentyl-2-phenylethyl, or 2-phenyl-2-pyridylethyl.
20. The compound of Claim 15, wherein R₉ is substituted or unsubstituted phenyl, substituted or unsubstituted phenyl-C₁-C₄-alkyl, diphenyl-C₁-C₄-alkyl, phenylfuranyl or heteroaryl-C₁-C₄-alkyl.
21. The compound of Claim 20, wherein R₉ is phenyl, phenyl-C₁-C₄-alkyl, diphenyl-C₁-C₄-alkyl wherein the phenyl group or phenyl groups bear one or more substituents independently selected from the group consisting of cyano, C₁-C₄-alkyl-S-, a halogen, a halogenated C₁-C₄-alkyl, C₁-C₄-alkoxy, trifluoromethyl, and substituted and unsubstituted phenoxy.
22. The compound of Claim 20, wherein R₉ is phenyl, phenyl-C₁-C₄-alkyl, diphenyl-C₁-C₄-alkyl wherein the phenyl group or phenyl groups bear one or more substituents independently selected from the group consisting of

cyano, methyl, methoxy, phenoxy, chloro-substituted phenoxy, methoxy-substituted phenoxy and methyl-substituted phenoxy.

23. The compound of Claim 15, wherein R_9 is phenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, diphenylmethyl, pyrazolylmethyl, 2,4-dimethylphenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-methyl-4-methoxyphenyl, 3-methyl-4-methoxyphenyl, 4-methylthiophenyl, 3-chlorophenyl, 3-trifluoromethylphenyl, benzyl, 2-trifluoromethylbenzyl, 3-trifluoromethylbenzyl, 2-chlorobenzyl, 3-chlorobenzyl, 4-chlorobenzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-fluorobenzyl, 3-azidylphenyl, 3-(4-methoxyphenoxy)phenyl, or 5-phenylfuran-2-yl.
24. The compound of Claim 15, wherein R_{10} is substituted or unsubstituted phenyl, alkyl substituted with a heteroaryl group, alkyl substituted with a heterocycloalkyl group, or an alkyl substituted with $-NR_{13}R_{14}$, wherein:
 R_{13} and R_{14} are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl; or
 R_{13} and R_{14} together with the nitrogen to which they are attached are a heterocycloalkyl.
25. The compound of Claim 24, wherein R_{10} is 2-(imidazol-4-yl)ethyl, 3-(imidazol-4-yl)propyl, 3-(imidazol-1-yl)propyl 2-(3-methylimidazol-4-yl)ethyl, 2-(morpholin-4-yl)ethyl, 2-(4-pyrazolyl)ethyl, 4-pyrazolylmethyl, 2-N,N-dimethylaminoethyl, 3-N,N-dimethylaminopropyl, or 2-(aminocarbonyl)phenyl.
26. The compound of Claim 15, wherein R_{11} is a linear or branched C_1 - C_4 -alkyl, substituted or unsubstituted phenyl, substituted or unsubstituted benzophenonyl, pyrazolyl, aminopyrazolyl, substituted or unsubstituted


indolyl-C₁-C₄-alkyl, thiophenyl, quinoxaline, substituted or unsubstituted phenyl-C₁-C₄-alkyl, pyridylcarbonylphenyl, phenylcarbonyl-C₁-C₄-alkyl, naphthyl, naphthyl-C₁-C₄-alkyl, diphenyl-C₁-C₄-alkyl, C₃-C₈-cycloalkyl-C₁-C₄-alkyl, C₁-C₄-alkylcarbonyl-C₁-C₄-alkyl, fluorenyl, pyrrolyl, N-methylpyrrolyl, or pyridyl.

27. The compound of Claim 26, wherein R₁₁ is a phenyl, phenyl-C₁-C₄-alkyl, phenylcarbonyl-C₁-C₄-alkyl, naphthyl-C₁-C₄-alkyl, diphenyl-C₁-C₄-alkyl, C₃-C₈-cycloalkyl-C₁-C₄-alkyl, fluorenyl or pyridyl substituted with one or more substituents independently selected from C₁-C₄-alkyl and C₁-C₄-alkoxy.
- 10 28. The compound of Claim 26, wherein R₁₁ is a benzophenonyl group, wherein said benzophenonyl group is substituted with a C₁-C₄-alkoxy group, a C₁-C₄-alkyl group or a chlorine atom.
29. The compound of Claim 15, wherein R₁₁ is benzophenon-2-yl, 4'-methoxybenzophenon-2-yl, 4'-chlorobenzophenon-2-yl, 2-(furan-2-yl)phenyl, 2-(thiophen-2-yl)phenyl, 2-benzylphenyl, 2-pyridylcarbonylphenyl, 2-(phenoxymethyl)phenyl, 2-(*t*-butylcarbonyl)phenyl, 2,2-diphenylethyl, 1-fluorenyl, (naphth-2-yl)methyl, naphth-1-yl, 3-(phenylcarbonyl)propyl, 4-phenylbutyl, 4-butylphenyl, 2-(4-chlorophenylcarbonyl)phenyl, 3-methoxyphenyl, N-methylpyrrol-2-yl, 2,3-dimethoxyphenyl, 3-butyl-2-pyridyl, 2-naphthylmethyl, 2-cyclohexylethyl, 3-methoxyphenyl, N-methyl-2-pyrrolyl, 2-cyclopentylethyl, 3-oxobutyl, 2-benzopyrazyl, quinoxalin-2-yl, 3-idolyl, (2-methylindol-3-yl)methyl, 3-(indol-3-yl)propyl, (indol-3-yl)methyl, (5-bromoindol-3-yl)methyl, 3-chlorophenyl, 3-aminopyrazol-4-yl, 2-(indol-3-yl)-1-hydroxyethyl, 3-fluorophenyl, 1-phenyl-1-hydroxymethyl, 2-phenylphenyl, 2-phenoxyphenyl, thiophen-2-yl, or isopropyl.
30. A composition of matter comprising an enantiomeric mixture of a compound represented by the following structural formula:




31. A compound which has a positive specific rotation, wherein the compound is represented by the following structural formula:


-87-



32. A compound which has a negative specific rotation, wherein the compound is represented by the following structural formula:



33. A method of treating a TNF- α mediated condition in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula I,




- 5 and physiological salts thereof, wherein:
 n is 0, 1 or 2;
 X is O, CH₂, S or SO₂;
 R₁ is H or NH₂;
 R₂ and R₃ are each, independently, -H, -OH, a substituted or
 10 unsubstituted alkyl, or a substituted or unsubstituted alkoxy;
 R₄ is, -H or a substituted or unsubstituted alkyl;
 V, W and Z are each, independently, N or CH; and
 Y is selected from the group consisting of substituted and
 unsubstituted phenyl, and substituted and unsubstituted heterocyclyl.
- 15 34. The method of Claim 33, wherein Y is a phenyl group which has one or more substituents independently selected from the group consisting of halogen, linear or branched C₁-C₄-alkoxy, trifluoromethoxy, dioxymethylene,

hydroxyalkyl, trifluoromethyl, HC(O)-, linear or branched C₁-C₄-alkyl, heterocyclyl and substituted or unsubstituted heterocycloalkylalkyl.

35. The method of Claim 34, wherein Y is a phenyl group which has one or more substituents selected from the group consisting of fluoro, chloro, methoxy, morpholyl, N-morpholinomethyl, tetrahydroisoquinolyl, tetrahydroisoquinolinomethyl, 4-(4-benzyl-piperazin-1-yl)methyl, 4-(4-(2-fluoro-phenyl)piperazin-1-yl)methyl, and isopropyl.
36. The method of Claim 33, wherein Y is selected from the group consisting of pyridyl, furyl, and pyrrolidyl.
37. The method of Claim 33, wherein the TNF- α mediated condition is selected from the group consisting of acute and chronic immune and autoimmune pathologies.
38. The method of Claim 37, wherein the TNF- α mediated condition is selected from the group consisting of systemic lupus erythematosus, rheumatoid arthritis, thyroidosis, graft versus host disease, scleroderma, diabetes mellitus and Graves' disease.
39. The method of Claim 33, wherein the TNF- α mediated condition is an infection.
40. The method of Claim 39, wherein the TNF- α mediated condition is selected from the group consisting of sepsis syndrome, cachexia, circulatory collapse and shock resulting from acute or chronic bacterial infection, acute and chronic parasitic, bacterial, viral and fungal infectious diseases.
41. The method of Claim 33, wherein the TNF- α mediated condition is an inflammatory disease.


42. The method of Claim 41, wherein the TNF- α mediated condition is selected from the group consisting of chronic inflammatory pathologies and vascular inflammatory pathologies.
43. The method of Claim 42, wherein the TNF- α mediated condition is selected from the group consisting of sarcoidosis, chronic inflammatory bowel disease, ulcerative colitis, Crohn's disease, disseminated intravascular coagulation, atherosclerosis, and Kawasaki's pathology.
44. The method of Claim 33, wherein the TNF- α mediated condition is a neurodegenerative disease.
45. The method of Claim 44, wherein the TNF- α mediated condition is selected from the group consisting of multiple sclerosis, acute transverse myelitis, lesions of the corticospinal system, disorders of the basal ganglia or cerebellar disorders, hyperkinetic movement disorders such as Huntington's Chorea and senile chorea, drug-induced movement disorders, hypokinetic movement disorders, progressive supranucleo palsy, astructural lesions of the cerebellum, spinal ataxia, Friedreich's ataxia, cerebellar cortical degenerations, multiple systems degenerations, Refsum's disease, abetalipoprotemia, ataxia, telangiectasia, mitochondrial multisystem disorder, multiple sclerosis, acute transverse myelitis, neurogenic muscular atrophies, Alzheimer's disease, Down's Syndrome in middle age, Diffuse Lewy body disease, Senile Dementia of Lewy body type, Wernicke-Korsakoff syndrome, chronic alcoholism, Creutzfeldt-Jakob disease, Subacute sclerosing panencephalitis, Hallerrorden-Spatz disease, and Dementia pugilistica.
46. The method of Claim 33, wherein the TNF- α mediated condition is cancer.

47. The method of Claim 46, wherein the TNF- α mediated condition is selected from the group consisting of TNF- α secreting tumors, leukemias, and lymphomas.
48. The method of Claim 33, wherein the TNF- α mediated condition is alcohol-induced hepatitis.
49. A method of treating a TNF- α mediated condition in a patient, comprising the step of administering to the patient a therapeutically effective amount of a compound represented by the following structural formula:




50. A method of treating a TNF- α mediated condition in a patient, comprising the step of administering to the patient a therapeutically effective amount of a compound represented by the following structural formula:

-92-



51. A method of treating a $\text{TNF-}\alpha$ mediated condition in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula II,

5



-93-

and physiological salts thereof, wherein,

R₅ is substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted cycloalkylalkyl;

R₆ is -H or -NR₁₃R₁₄;

5 R₇ is substituted or unsubstituted phenyl; and

R₁₃ and R₁₄ are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl; or

10 R₁₃ and R₁₄ together with the nitrogen to which they are attached are a heterocycloalkyl.

52. The method of Claim 51, wherein R₅ is substituted or unsubstituted benzyl.
53. The method of Claim 52, wherein R₅ is benzyl having one or more substituents independently selected from the group consisting of halogen, linear C₁-C₄-alkoxy and branched C₁-C₄-alkoxy.
- 15 54. The method of Claim 53, wherein R₅ is benzyl having one or more substituents independently selected from the group consisting of chloro and methoxy.
55. The method of Claim 51, wherein R₅ is C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-C₁-C₄-alkyl or substituted or unsubstituted phenyl-C₂-C₄-alkyl.
- 20 56. The method of Claim 54, wherein R₅ is selected from the group consisting of 2-phenethyl, cyclohexyl and cyclopentylethyl.
57. The method of Claim 51, wherein R₇ is phenyl having one or more substituents independently selected from the group consisting of halogen, linear C₁-C₆-alkyl, branched C₁-C₆-alkyl and cyclic C₃-C₆-alkyl and
- 25 trifluoromethyl.


58. The method of Claim 57, wherein R₇ is phenyl having one or more substituents independently selected from the group consisting of fluoro, chloro, linear C₁-C₄-alkyl, and branched C₁-C₄-alkyl.
59. The method of Claim 51, wherein the TNF- α mediated condition is selected from the group consisting of acute and chronic immune and autoimmune pathologies.
60. The method of Claim 59, wherein the TNF- α mediated condition is selected from the group consisting of systemic lupus erythematosus, rheumatoid arthritis, thyroidosis, graft versus host disease, scleroderma, diabetes mellitus and Graves' disease.
61. The method of Claim 51, wherein the TNF- α mediated condition is an infection.
62. The method of Claim 61, wherein the TNF- α mediated condition is selected from the group consisting of sepsis syndrome, cachexia, circulatory collapse and shock resulting from acute or chronic bacterial infection, acute and chronic parasitic, bacterial, viral and fungal infectious diseases.
63. The method of Claim 51, wherein the TNF- α mediated condition is an inflammatory disease.
64. The method of Claim 63, wherein the TNF- α mediated condition is selected from the group consisting of chronic inflammatory pathologies and vascular inflammatory pathologies.
65. The method of Claim 64, wherein the TNF- α mediated condition is selected from the group consisting of sarcoidosis, chronic inflammatory bowel disease, ulcerative colitis, Crohn's disease, disseminated intravascular

-95-

coagulation, atherosclerosis, and Kawasaki's pathology.

66. The method of Claim 51, wherein the TNF- α mediated condition is a neurodegenerative disease.
67. The method of Claim 66, wherein the TNF- α mediated condition is selected from the group consisting of multiple sclerosis, acute transverse myelitis, lesions of the corticospinal system, disorders of the basal ganglia or cerebellar disorders, hyperkinetic movement disorders such as Huntington's Chorea and senile chorea, drug-induced movement disorders, hypokinetic movement disorders, progressive supranucleo palsy, astructural lesions of the cerebellum, spinal ataxia, Friedreich's ataxia, cerebellar cortical degenerations, multiple systems degenerations, Refsum's disease, abetalipoproteemia, ataxia, telangiectasia, mitochondrial multisystem disorder, multiple sclerosis, acute transverse myelitis, neurogenic muscular atrophies, Alzheimer's disease, Down's Syndrome in middle age, Diffuse Lewy body disease, Senile Dementia of Lewy body type, Wernicke-Korsakoff syndrome, chronic alcoholism, Creutzfeldt-Jakob disease, Subacute sclerosing panencephalitis, Hallerrorden-Spatz disease, and Dementia pugilistica.
68. The method of Claim 51, wherein the TNF- α mediated condition is cancer.
69. The method of Claim 68, wherein the TNF- α mediated condition is selected from the group consisting of TNF- α secreting tumors, leukemias, and lymphomas.
70. The method of Claim 51, wherein the TNF- α mediated condition is alcohol-induced hepatitis.

71. A method of treating a TNF- α mediated condition in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula III,



5 and physiological salts thereof, wherein:

R_8 and R_{12} are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, substituted or unsubstituted aralkyl or a substituted or unsubstituted heteroaralkyl;

10 R_9 is -H, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl, a substituted or unsubstituted heteroaryl or a substituted or unsubstituted heteroaralkyl;

R_{10} is substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaralkyl, or a substituted or unsubstituted heterocycloalkylalkyl; and

15 R_{11} is a substituted or unsubstituted alkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted aralkyl, a substituted or unsubstituted cycloalkylalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted heteroaralkyl, a substituted or unsubstituted benzophenonyl, or a substituted or unsubstituted cycloalkylalkyl.

-97-

72. The method of Claim 71, wherein one of R_8 or R_{12} is -H and the other is substituted or unsubstituted phenyl, phenyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -alkyl, linear C_1 - C_{12} -alkyl, branched C_1 - C_{12} -alkyl, cyclic C_3 - C_{12} -alkyl, or dicycloalkyl- C_1 - C_4 -alkyl.
- 5 73. The method of Claim 72, wherein one of R_8 or R_{12} is -H and the other is phenyl, phenyl- C_1 - C_4 -alkyl, or diphenyl- C_1 - C_4 -alkyl wherein the phenyl group or phenyl groups bear one or more substituents independently selected from the group consisting of C_1 - C_4 -alkoxy, C_1 - C_4 -alkyl and cyano.
- 10 74. The method of Claim 73, wherein the phenyl group or phenyl groups bear one or more substituents independently selected from the group consisting of methoxy, methyl and cyano.
- 15 75. The method of Claim 71, wherein R_8 is selected from the group consisting of 2,2-diphenylethyl, 2-(4-ethylphenyl)ethyl, benzyl, diphenylmethyl, 1,2-diphenylethyl, 3,3-diphenylpropyl, 3,4,5-trimethoxybenzyl, 2,4,4-trimethylisopentyl, 2-(4-methoxyphenyl)ethyl, 2-cyclopentyl-2-phenylethyl, or 2-phenyl-2-pyridylethyl.
- 20 76. The method of Claim 71 wherein R_9 is substituted or unsubstituted phenyl, phenyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -alkyl, phenylfuranyl or heteroaryl- C_1 - C_4 -alkyl.
- 25 77. The method of Claim 76, wherein R_9 is phenyl, phenyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -alkyl wherein the phenyl group or phenyl groups bear one or more substituents independently selected from the group consisting of cyano, C_1 - C_4 -alkyl-S-, a halogen C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, trifluoromethyl, and substituted and unsubstituted phenoxy.

78. The method of Claim 76, wherein R_9 is phenyl, phenyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -alkyl wherein the phenyl group or phenyl groups bear one or more substituents independently selected from the group consisting of cyano, methyl, methoxy, phenoxy, chloro-substituted phenoxy, methoxy-substituted phenoxy and methyl-substituted phenoxy.
79. The compound of Claim 71, wherein R_9 is phenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, diphenylmethyl, pyrazolylmethyl, 2,4-dimethylphenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-methyl-4-methoxyphenyl, 3-methyl-4-methoxyphenyl, 4-methylthiophenyl, 3-chlorophenyl, 3-trifluoromethylphenyl, benzyl, 2-trifluoromethylbenzyl, 3-trifluoromethylbenzyl, 2-chlorobenzyl, 3-chlorobenzyl, 4-chlorobenzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-fluorobenzyl, 3-azidylphenyl, 3-(4-methoxyphenoxy)phenyl, or 5-phenylfuran-2-yl.
80. The method of Claim 71, wherein R_{10} is substituted or unsubstituted phenyl, alkyl substituted with a heteroaryl group, alkyl substituted with a heterocycloalkyl group, or an alkyl substituted with $-NR_{13}R_{14}$, wherein:
- R_{13} and R_{14} are each, independently, -H, a substituted or unsubstituted alkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted aryl, or a substituted or unsubstituted aralkyl; or
- R_{13} and R_{14} together with the nitrogen to which they are attached are a heterocycloalkyl.
81. The method of Claim 80, wherein R_{10} is 2-(imidazol-4-yl)ethyl, 3-(imidazol-4-yl)propyl, 3-(imidazol-1-yl)propyl 2-(3-methylimidazol-4-yl)ethyl, 2-(morpholin-4-yl)ethyl, 2-(4-pyrazolyl)ethyl, 4-pyrazolylmethyl, 2-N,N-dimethylaminoethyl, 3-N,N-dimethylaminopropyl, and 2-(aminocarbonyl)phenyl.

82. The method of Claim 71, wherein R_{11} is a linear or branched C_1 - C_4 -alkyl, substituted or unsubstituted phenyl, substituted or unsubstituted benzophenonyl, pyrazolyl, aminopyrazolyl, substituted or unsubstituted indolyl- C_1 - C_4 -alkyl, thiophenyl, quinoxaline, substituted or unsubstituted phenyl- C_1 - C_4 -alkyl, pyridylcarbonylphenyl, phenylcarbonyl- C_1 - C_4 -alkyl, naphthyl, naphthyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -alkyl, C_5 - C_8 -cycloalkyl- C_1 - C_4 -alkyl, C_1 - C_4 -alkylcarbonyl- C_1 - C_4 -alkyl, fluorenyl, pyrrolyl, N-methylpyrrolyl, or pyridyl.
83. The method of Claim 82, wherein R_{11} is a phenyl, phenyl- C_1 - C_4 -alkyl, phenylcarbonyl- C_1 - C_4 -alkyl, naphthyl- C_1 - C_4 -alkyl, diphenyl- C_1 - C_4 -alkyl, C_5 - C_8 -cycloalkyl- C_1 - C_4 -alkyl, fluorenyl or pyridyl substituted with one or more substituents independently selected from C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy.
84. The method of Claim 82, wherein R_{11} is a benzophenonyl group, wherein said benzophenonyl group is substituted with a C_1 - C_4 -alkoxy group, a C_1 - C_4 -alkyl group or a chlorine atom.
85. The method of Claim 71, wherein R_{11} is benzophenon-2-yl, 4'-methoxybenzophenon-2-yl, 4'-chlorobenzophenon-2-yl, 2-(furan-2-yl)phenyl, 2-(thiophen-2-yl)phenyl, 2-benzylphenyl, 2-pyridylcarbonylphenyl, 2-(phenoxymethyl)phenyl, 2-(*t*-butylcarbonyl)phenyl, 2,2-diphenylethyl, 1-fluorenyl, (naphth-2-yl)methyl, naphth-1-yl, 3-(phenylcarbonyl)propyl, 4-phenylbutyl, 4-butylphenyl, 2-(4-chlorophenylcarbonyl)phenyl, 3-methoxyphenyl, N-methylpyrrol-2-yl, 2,3-dimethoxyphenyl, 3-butyl-2-pyridyl, 2-naphthylmethyl, 2-cyclohexylethyl, 3-methoxyphenyl, N-methyl-2-pyrrolyl, 2-cyclopentylethyl, 3-oxobutyl, 2-benzopyrazyl, quinoxalin-2-yl, 3-idolyl, (2-methylindol-3-yl)methyl, 3-(indol-3-yl)propyl, (indol-3-yl)methyl, (5-bromoindol-3-yl)methyl, 3-chlorophenyl, 3-aminopyrazol-4-yl, 2-(indol-3-yl)-1-hydroxyethyl, 3-fluorophenyl, 1-phenyl-1-hydroxymethyl, 2-phenylphenyl, 2-phenoxyphenyl, thiophen-2-yl, or isopropyl.


-100-

86. The method of Claim 71, wherein the TNF- α mediated condition is selected from the group consisting of acute and chronic immune and autoimmune pathologies.
- 5 87. The method of Claim 86, wherein the TNF- α mediated condition is selected from the group consisting of systemic lupus erythematosus, rheumatoid arthritis, thyroidosis, graft versus host disease, scleroderma, diabetes mellitus and Graves' disease.
88. The method of Claim 71, wherein the TNF- α mediated condition is an infection.
- 10 89. The method of Claim 88, wherein the TNF- α mediated condition is selected from the group consisting of sepsis syndrome, cachexia, circulatory collapse and shock resulting from acute or chronic bacterial infection, acute and chronic parasitic, bacterial, viral and fungal infectious diseases.
90. The method of Claim 71, wherein the TNF- α mediated condition is an
15 inflammatory disease.
91. The method of Claim 90 wherein the TNF- α mediated condition is selected from the group consisting of chronic inflammatory pathologies and vascular inflammatory pathologies.
- 20 92. The method of Claim 91, wherein the TNF- α mediated condition is selected from the group consisting of sarcoidosis, chronic inflammatory bowel disease, ulcerative colitis, Crohn's disease, disseminated intravascular coagulation, atherosclerosis, and Kawasaki's pathology.

-101-

93. The method of Claim 71, wherein the TNF- α mediated condition is a neurodegenerative disease.
94. The method of Claim 93, wherein the TNF- α mediated condition is selected from the group consisting of multiple sclerosis, acute transverse myelitis, lesions of the corticospinal system, disorders of the basal ganglia or cerebellar disorders, hyperkinetic movement disorders such as Huntington's Chorea and senile chorea, drug-induced movement disorders, hypokinetic movement disorders, progressive supranucleo palsy, astructural lesions of the cerebellum, spinal ataxia, Friedreich's ataxia, cerebellar cortical degenerations, multiple systems degenerations, Refsum's disease, abetalipoproteemia, ataxia, telangiectasia, mitochondrial multisystem disorder, multiple sclerosis, acute transverse myelitis, neurogenic muscular atrophies, Alzheimer's disease, Down's Syndrome in middle age, Diffuse Lewy body disease, Senile Dementia of Lewy body type, Wernicke-Korsakoff syndrome, chronic alcoholism, Creutzfeldt-Jakob disease, Subacute sclerosing panencephalitis, Hallerrorden-Spatz disease, and Dementia pugilistica.
95. The method of Claim 71, wherein the TNF- α mediated condition is cancer.
96. The method of Claim 95, wherein the TNF- α mediated condition is selected from the group consisting of TNF- α secreting tumors, leukemias, and lymphomas.
97. The method of Claim 71, wherein the TNF- α mediated condition is alcohol-induced hepatitis.
98. A method of treating a TNF-a mediated condition in a patient, comprising the step of administering to the patient a therapeutically effective amount of a compound represented by the following structural formula:

-102-




99. The method of Claim 98, wherein the compound has a positive specific rotation.

100. The method of Claim 98, wherein the compound has a negative specific rotation.


101. A method of treating multiple sclerosis in a patient, comprising the step of administering to the patient a therapeutically effective amount of a compound represented by the following structural formula:

5

-103-



102. A compound of Formula I,



and physiological salts thereof, wherein:

n is 0, 1 or 2;

X is O, CH₂, S or SO₂;

-104-

R_1 is H or NH_2 ;


R_2 and R_3 are each, independently, -H, -OH, a substituted or unsubstituted alkyl, or a substituted or unsubstituted alkoxy;

R_4 is, -H or a substituted or unsubstituted alkyl;

5

V, W and Z are each, independently, N or CH; and

Y is represented by the following structural formula:




10 wherein R_{50} and R_{51} are independently an alkyl group, a substituted alkyl group, an aryl group a substituted aryl group, or, taken together with the nitrogen atom to which they are bonded, are a substituted heterocycloalkyl, an unsubstituted heterocycloalkyl, a substituted heteroaryl group or an unsubstituted heteroaryl group.

103. A method of treating a $TNF-\alpha$ mediated condition in a patient, comprising administering to the patient a therapeutically effective amount of the compound of Claim 102.

15

104. A compound represented by the following structural formula:


-105-



and physiologically acceptable salts thereof.

105. A method of treating a $\text{TNF-}\alpha$ mediated condition in a patient, comprising administering to the patient a therapeutically effective amount of the compound of Claim 104.
- 5
106. A compound represented by the following structural formula:

-106-



and physiologically acceptable salts thereof.

107. A method of treating a TNF- α mediated condition in a patient, comprising administering to the patient a therapeutically effective amount of the compound of Claim 106.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 November 2001 (22.11.2001)

PCT

(10) International Publication Number
WO 01/87849 A3

(51) International Patent Classification⁷: **C07D 233/54**,
A61K 31/395, A61P 43/00, C07D 401/04, 401/14, 405/12,
409/12, 403/12, 403/14

(74) Agents: **BROOK, David, E. et al.**; Hamilton, Brook,
Smith & Reynolds, P.C., 530 Virginia Road, P.O. Box
9133, Concord, MA 01742-9133 (US).

(21) International Application Number: **PCT/US01/15027**

(22) International Filing Date: **10 May 2001 (10.05.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/203,784 12 May 2000 (12.05.2000) US
60/205,213 18 May 2000 (18.05.2000) US

(71) Applicant (for all designated States except US): **GEN-
ZYME CORPORATION** [US/US]; One Kendall Square,
Cambridge, MA 02139 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SNEDDON, Scott**,
F. [US/US]; 38 Washington Square South, Salem, MA
01970 (US). **KANE, John, L.** [US/US]; 10 O'Moore
Avenue, Maynard, MA 01754 (US). **HIRTH, Bradford**,
H. [US/US]; 4 Dogwood Road, Littleton, MA 01460 (US).
VINICK, Fred [US/US]; 5 Litchfield Road, Lexington,
MA 02420 (US). **QIAO, Shuang** [CN/US]; 334 Harvard
Street, Apt. J-3, Cambridge, MA 02139 (US). **NAHILL**,
Sharon, R. [US/US]; 21 Vincent Avenue, Belmont, MA
02478 (US).


Published:

— with international search report


(88) Date of publication of the international search report:
6 June 2002

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.


(54) Title: **MODULATORS OF TNF- ALPHA SIGNALING**



(I)



(II)



(III)

(57) Abstract: The present invention provides compounds which are modulators of TNF- α signaling and methods of use thereof for treating a patient having a TNF- α mediated condition. The compounds can be represented by structural formulas (I), (II) and (III).

WO 01/87849 A3

INTERNATIONAL SEARCH REPORT

Int lional Application No

PCT/US 01/15027

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D233/54 A61K31/395 A61P43/00 C07D401/04 C07D401/14
C07D405/12 C07D409/12 C07D403/12 C07D403/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JAY WROBEL ET AL.: "Syntheses of Tolrestat analogues containing additional substituents in the ring and their evaluation as aldose reductase inhibitors. Identification of potent, orally active 2-fluoro derivatives" JOURNAL OF MEDICINAL CHEMISTRY., vol. 34, no. 8, - 1991 pages 2504-2520, XP000567039 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 * table 1, compounds 8a-8f * --- -/--	15



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

22 January 2002

Date of mailing of the international search report

08/02/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Int lional Application No

PCT/US 01/15027

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>CHEMICAL ABSTRACTS, vol. 107, no. 5, 3 August 1987 (1987-08-03) Columbus, Ohio, US; abstract no. 32462w, COUTTS, R. T. ET AL.: "Acetylation and pentafluorobenzoylation of lidocaine metabolites in aqueous solution and identidication of derivatives by combined gas chromatography/mass spectrometry." XP002188016 abstract & BIOMED. ENVIRON. MASS SPECTROM., vol. 14, no. 4, - 1987 pages 173-182, -& DATABASE CHEMICAL ABSTRACTS 'Online! CA 107:32462, XP002188018 compound with RN 108966-34-3 and -28-5</p>	15
X	<p>JORDI GARCIA ET AL.: "New synthetic "tricks". Triphenylphosphine-mediated amide formation from carboxylic acids and azides" TETRAHEDRON LETTERS., vol. 25, no. 42, - 1984 pages 4841-4844, XP002188015 ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM., NL ISSN: 0040-4039 * page 4842: product of third additional example *</p>	15
X	<p>CHEMICAL ABSTRACTS, vol. 130, no. 11, 15 March 1999 (1999-03-15) Columbus, Ohio, US; abstract no. 139333g, KONDO, KAZUMI ET AL.: "Preparation of amides as vasopressin antagonists, oxytocin antagonists, and vasopressin agonists." XP002188017 abstract & JP 01 101456 A (OHTSUKA PHARMACEUTICAL CO., LTD.) 19 April 1989 (1989-04-19) -& DATABASE CHEMICAL ABSTRACTS 'Online! CA 130: 139333, XP002188019 compound with RN 219988-64-4, -72-4, -73-5, -79-1 and -80-4</p>	15
X	<p>DE 29 37 698 A (A. NATTERMANN & CIE GMBH) 2 April 1981 (1981-04-02) examples 1-5</p>	15

-/-

INTERNATIONAL SEARCH REPORT

Int l tional Application No

PCT/US 01/15027

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 26209 A (NOVARTIS AG) 11 May 2000 (2000-05-11) claims -----	1
A	WO 99 00363 A (PHARMACIA & UPJOHN S.P.A.) 7 January 1999 (1999-01-07) claims -----	1
A	US 6 048 841 A (ANDREW DOUGLAS BAXTER ET AL.) 11 April 2000 (2000-04-11) claims -----	15

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Claims Nos.: 15-29 (partially)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty of compounds of formula III of claim 15. So many documents were retrieved that it is impossible to determine which parts of the claim may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of claim 15 is impossible. Consequently, with respect to claim 15 and its dependent claims, the search and the search report can only be considered comprehensive for the compounds of the specific examples of table I of the present application.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/15027

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 1101456	A	19-04-1989	NONE	
DE 2937698	A	02-04-1981	DE 2937698 A1	02-04-1981
			BE 885268 A1	16-01-1981
			FR 2465712 A1	27-03-1981
			GB 2058752 A ,B	15-04-1981
			IT 1131960 B	25-06-1986
			JP 1410815 C	24-11-1987
			JP 56059743 A	23-05-1981
			JP 62012788 B	20-03-1987
			US 4303673 A	01-12-1981
WO 0026209	A	11-05-2000	AU 6476599 A	22-05-2000
			WO 0026209 A1	11-05-2000
WO 9900363	A	07-01-1999	AU 8727498 A	19-01-1999
			WO 9900363 A1	07-01-1999
US 6048841	A	11-04-2000	AU 711907 B2	21-10-1999
			AU 7634396 A	11-06-1997
			BR 9611636 A	28-12-1999
			CN 1202895 A ,B	23-12-1998
			CZ 9801455 A3	12-08-1998
			EP 0874842 A1	04-11-1998
			WO 9719075 A1	29-05-1997
			HU 9904131 A2	28-06-2000
			JP 2000500482 T	18-01-2000
			NO 982290 A	20-05-1998
			PL 326710 A1	26-10-1998